

005412

B-JGT-K-NBA

MATHEMATICS

Paper I

Time Allowed : Three Hours

Maximum Marks : 200

INSTRUCTIONS

Candidates should attempt questions 1 and 5 which are compulsory, and any THREE of the remaining questions, selecting at least ONE question from each Section.

All questions carry equal marks.

Marks allotted to parts of a question are indicated against each.

Answers must be written in ENGLISH only.

Assume suitable data, if considered necessary, and indicate the same clearly.

Unless indicated otherwise, symbols & notations carry their usual meaning.

SECTION A

1. Answer any *five* of the following :

(a) Show that the set

$$P[t] = \{at^2 + bt + c \mid a, b, c \in \mathbb{R}\}$$

forms a vector space over the field \mathbb{R} . Find a basis for this vector space. What is the dimension of this vector space ?

8

B-JGT-K-NBA

1

[Contd.]

- (b) Determine whether the quadratic form

$$q = x^2 + y^2 + 2xz + 4yz + 3z^2$$

is positive definite.

8

- (c) Prove that between any two real roots of $e^x \sin x = 1$, there is at least one real root of $e^x \cos x + 1 = 0$.

8

- (d) Let f be a function defined on \mathbb{R} such that

$$f(x + y) = f(x) + f(y), \quad x, y \in \mathbb{R}.$$

If f is differentiable at one point of \mathbb{R} , then prove that f is differentiable on \mathbb{R} .

8

- (e) If a plane cuts the axes in A, B, C and (a, b, c) are the coordinates of the centroid of the triangle ABC , then show that the equation of the plane is

$$\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 3.$$

8

- (f) Find the equations of the spheres passing through the circle

$$x^2 + y^2 + z^2 - 6x - 2z + 5 = 0, \quad y = 0$$

and touching the plane $3y + 4z + 5 = 0$.

8

- (a) Show that the vectors

$$\alpha_1 = (1, 0, -1), \quad \alpha_2 = (1, 2, 1), \quad \alpha_3 = (0, -3, 2)$$

form a basis for \mathbb{R}^3 . Find the components of $(1, 0, 0)$ w.r.t. the basis $\{\alpha_1, \alpha_2, \alpha_3\}$.

10

(b) Find the characteristic polynomial of

$$\begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 2 \\ 0 & 1 & 3 \end{pmatrix}. \text{ Verify Cayley - Hamilton theorem}$$

for this matrix and hence find its inverse. 10

(c) Let $A = \begin{pmatrix} 5 & -6 & -6 \\ -1 & 4 & 2 \\ 3 & -6 & -4 \end{pmatrix}$. Find an invertible matrix P such that $P^{-1}AP$ is a diagonal matrix. 12

(d) Find the rank of the matrix

$$\begin{pmatrix} 1 & 2 & 1 & 1 & 2 \\ 2 & 4 & 3 & 4 & 7 \\ -1 & -2 & 2 & 5 & 3 \\ 3 & 6 & 2 & 1 & 3 \\ 4 & 8 & 6 & 8 & 9 \end{pmatrix}$$

8

(a) Discuss the convergence of the integral

$$\int_0^{\infty} \frac{dx}{1+x^4 \sin^2 x}$$

10

(b) Find the extreme value of xyz if $x + y + z = a$. 10

(c) Let

$$f(x, y) = \begin{cases} \frac{xy(x^2 - y^2)}{x^2 + y^2}, & \text{if } (x, y) \neq (0, 0) \\ 0, & \text{if } (x, y) = (0, 0) \end{cases}$$

Show that :

(i) $f_{xy}(0, 0) \neq f_{yx}(0, 0)$

(ii) f is differentiable at $(0, 0)$ 10

(d) Evaluate $\iint_D (x + 2y) \, dA$, where D is the region bounded by the parabolas $y = 2x^2$ and $y = 1 + x^2$. 10

4. (a) Prove that the second degree equation

$$x^2 - 2y^2 + 3z^2 + 5yz - 6zx - 4xy + 8x - 19y - 2z - 20 = 0$$

represents a cone whose vertex is $(1, -2, 3)$. 10

(b) If the feet of three normals drawn from a point P

to the ellipsoid $\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$ lie in the

plane $\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$, prove that the feet of the

other three normals lie in the plane

$$\frac{x}{a} + \frac{y}{b} + \frac{z}{c} + 1 = 0. \quad \text{10}$$

- (c) If $\frac{x}{1} = \frac{y}{2} = \frac{z}{3}$ represents one of the three mutually perpendicular generators of the cone $5yz - 8zx - 3xy = 0$, find the equations of the other two. 10

- (d) Prove that the locus of the point of intersection of three tangent planes to the ellipsoid

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1, \text{ which are parallel to the}$$

conjugate diametral planes of the ellipsoid

$$\frac{x^2}{\alpha^2} + \frac{y^2}{\beta^2} + \frac{z^2}{\gamma^2} = 1 \text{ is}$$

$$\frac{x^2}{\alpha^2} + \frac{y^2}{\beta^2} + \frac{z^2}{\gamma^2} = \frac{a^2}{\alpha^2} + \frac{b^2}{\beta^2} + \frac{c^2}{\gamma^2}$$

10

SECTION B

Answer any **five** of the following :

- (a) Show that $\cos(x + y)$ is an integrating factor of

$$y \, dx + [y + \tan(x + y)] \, dy = 0.$$

Hence solve it.

8

- (b) Solve

$$\frac{d^2y}{dx^2} - 2\frac{dy}{dx} + y = xe^x \sin x$$

8

- (c) A uniform rod AB rests with one end on a smooth vertical wall and the other on a smooth inclined plane, making an angle α with the horizon. Find the positions of equilibrium and discuss stability.

8

- (d) A particle is thrown over a triangle from one end of a horizontal base and grazing the vertex falls on the other end of the base. If θ_1 and θ_2 be the base angles and θ be the angle of projection, prove that,

$$\tan \theta = \tan \theta_1 + \tan \theta_2.$$

8

- (e) Prove that the horizontal line through the centre of pressure of a rectangle immersed in a liquid with one side in the surface, divides the rectangle in two parts, the fluid pressure on which, are in the ratio, 4 : 5.

8

- (f) Find the directional derivation of \vec{V}^2 , where,
 $\vec{V} = xy^2\vec{i} + zy^2\vec{j} + xz^2\vec{k}$ at the point (2, 0, 3)
 in the direction of the outward normal to the
 surface $x^2 + y^2 + z^2 = 14$ at the point (3, 2, 1). . 8

6. (a) Solve the following differential equation

$$\frac{dy}{dx} = \sin^2(x - y + 6) \quad 8$$

- (b) Find the general solution of.

$$\frac{d^2y}{dx^2} + 2x \frac{dy}{dx} + (x^2 + 1)y = 0 \quad 12$$

- (c) Solve

$$\left(\frac{d}{dx} - 1\right)^2 \left(\frac{d^2}{dx^2} + 1\right)^2 y = x + e^x \quad 10$$

- (d) Solve by the method of variation of parameters
 the following equation

$$(x^2 - 1) \frac{d^2y}{dx^2} - 2x \frac{dy}{dx} + 2y = (x^2 - 1)^2 \quad 10$$

7. (a) A uniform chain of length $2l$ and weight W , is
 suspended from two points A and B in the same
 horizontal line. A load P is now hung from the
 middle point D of the chain and the depth of
 this point below AB is found to be h . Show that
 each terminal tension is,

$$\frac{1}{2} \left[P \cdot \frac{l}{h} + W \cdot \frac{h^2 + l^2}{2hl} \right] \quad 14$$

- (b) A particle moves with a central acceleration $\frac{\mu}{(\text{distance})^2}$, it is projected with velocity V at a distance R . Show that its path is a rectangular hyperbola if the angle of projection is,

$$\sin^{-1} \left[\frac{\mu}{VR \left(V^2 - \frac{2\mu}{R} \right)^{1/2}} \right]$$

13

- (c) A smooth wedge of mass M is placed on a smooth horizontal plane and a particle of mass m slides down its slant face which is inclined at an angle α to the horizontal plane. Prove that the acceleration of the wedge is,

$$\frac{mg \sin \alpha \cos \alpha}{M + m \sin^2 \alpha}$$

13

8. (a) (i) Show that

$$\vec{F} = (2xy + z^3) \vec{i} + x^2 \vec{j} + 3z^2 x \vec{k}$$

is a conservative field. Find its scalar potential and also the work done in moving a particle from $(1, -2, 1)$ to $(3, 1, 4)$.

5

- (ii) Show that, $\nabla^2 f(r) = \left(\frac{2}{r} \right) f'(r) + f''(r)$, where

$$r = \sqrt{x^2 + y^2 + z^2}$$

5

- (b) Use divergence theorem to evaluate,

$$\iiint_S (x^3 \, dy \, dz + x^2 y \, dz \, dx + x^2 z \, dy \, dx),$$

where S is the sphere, $x^2 + y^2 + z^2 = 1$. 10

- (c) If $\vec{A} = 2y \vec{i} - z \vec{j} - x^2 \vec{k}$ and S is the surface of the parabolic cylinder $y^2 = 8x$ in the first octant bounded by the planes $y = 4$, $z = 6$, evaluate the surface integral,

$$\iint_S \vec{A} \cdot \hat{n} \, dS.$$

10

- (d) Use Green's theorem in a plane to evaluate the integral, $\int_C [(2x^2 - y^2) \, dx + (x^2 + y^2) \, dy]$, where C

is the boundary of the surface in the xy -plane enclosed by, $y = 0$ and the semi-circle,

$$y = \sqrt{1 - x^2}. \quad \text{10}$$