

# MATHEMATICS



## PAPER - I SECTION A

1. Answer any four of the following:

- (a) Let  $V$  be the vector space of polynomials in  $x$  with real coefficients of degree at most 2.

Let  $t_1, t_2, t_3$  be any 3 distinct real numbers. Define

$$L_i : V \rightarrow \mathbb{R} \text{ by } L_i(f) = f(t_i), i = 1, 2, 3. \text{ Show that}$$

- (i)  $L_1, L_2, L_3$  are linear functionals on  $V$   
 (ii)  $\{L_1, L_2, L_3\}$  is a basis for the dual space  $V^*$  of  $V$ .

(10)

- (b) In the notation of (a) above, find a basis  $B = \{p_1, p_2, p_3\}$  for  $V$  which is dual to  $\{L_1, L_2, L_3\}$  and also express each  $P \in V$  in terms of elements of  $B$ .

- (c) Let  $f$  be a function defined on  $[0, 1]$  by

$$f(x) = \begin{cases} 0, & \text{if } x \text{ is irrational} \\ \frac{1}{q}, & \text{if } x = \frac{p}{q}, q \neq 0 \end{cases}$$

and  $p, q$ , are relatively prime positive integers. Show that  $f$  is continuous at each irrational point and discontinuous at each rational point  $\frac{p}{q}$ .

(10)

- (d) Show that the function  $[x]$ , where  $[x]$  denotes the greatest integer not greater than  $x$ , is integrable in  $[0, 3]$ . Also evaluate  $\int_0^3 [x] dx$ .

(10)

- (e) Prove that the polar of one limiting point of a coaxial system of circles with respect to any circle of the system passes through the other limiting point.

(10)

2. (a) Let  $V$  be the vector space of polynomials in  $x$  with complex coefficients. Define

$$T: V \rightarrow V \text{ by } (Tf)(x) = xf(x) \text{ and}$$

$$U: V \rightarrow V \text{ by } U\left(\sum_{i=0}^n c_i x^i\right) = \sum_{i=0}^{n-1} c_i + 1x^i$$

Find (i)  $\ker T$  (ii) show  $U$  is linear (iii) show that  $UT = I$  and  $TU \neq I$ ,  $I$  = identity on  $V$ .

(10)

- (b) Let  $A = \begin{pmatrix} 1 & 0 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}$  show that for every integer  $n \geq 3$ ,  $A^n = A^{n-2} + A^2 - I$  and hence find the matrix  $A^8$ .

(10)

- (c) Find the characteristic and minimal polynomials of

$$A = \begin{pmatrix} 2 & 0 & 0 \\ 1 & 2 & 0 \\ 0 & 0 & -1 \end{pmatrix}$$

and determine whether A is diagonalizable.

(10)

(d) Let  $A = \begin{pmatrix} 1 & 1 & 2 \\ -1 & 2 & 1 \\ 0 & 1 & 3 \end{pmatrix}$

Find an invertible  $3 \times 3$  matrix P such that  $P^{-1}AP = D$ , D is diagonal matrix. Find D also.

(10)

3. (a) Examine the convergence of the integral

$$\int_0^1 x^{n-1} \log x dx.$$

(10)

- (b) Examine the function

$$f(x, y) = \begin{cases} \frac{xy}{\sqrt{x^2 + y^2}} & (x, y) \neq (0, 0) \\ 0 & (x, y) = (0, 0) \end{cases}$$

For continuity, partial derivability of the first order and differentiability at (0, 0).

(10)

- (c) Find the maximum and minimum values of the function  $f(x, y, z) = xy + 2z$  on the circle which is the intersection of the plane  $x + y + z = 0$  and the sphere  $x^2 + y^2 + z^2 = 24$ .

(10)

- (d) Find the volume of there region R lying below the plane  $z = 3 + 2y$  and above the paraboloid  $z = x^2 + y^2$ .

(10)

4. (a) CP and CD are conjugate diameters of an ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ . Prove that the locus of the orthocenter of the triangle CPD is the curve  $2(b^2y^2 + a^2x^2)^3 = (a^2 - b^2)^2(b^2y^2 - a^2x^2)^2$ .

(13)

- (b) If  $\frac{x}{1} = \frac{y}{2} = \frac{z}{3}$  represents one of the three mutually perpendicular generators of the cone  $5yz - 8zx - 3xy = 0$  Find the equations of the other two.

(13)

- (v) If the section of the enveloping cone of the ellipsoid

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1,$$

Whose vertex is P, by the plane  $z = 0$  is a rectangular hyperbola, prove that the locus of P is

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} - \frac{z^2}{c^2} = 1.$$

(14)

## SECTION B

5. Answer any four of the following:

- (a) A tank of 100 liters capacity is initially full of water. Pure water is allowed to run into the tank at the rate of 1 liter per minute and at the same time salt water containing  $\frac{1}{4}$  kg of salt per liter flows into the tank at the rate of 1 liter per minute. The mixture (there is perfect mixing in the tank at all times) flows out at the rate of 2 liters per minute. Form the differential equation and find the amount of salt in the tank after  $t$  minutes. Find this when  $t = 50$  minutes.

(10)

- (b) A constant coefficient differential equation has auxiliary equation expressible in factored form as

$$P(m) = m^3 (m - 1)^2 (m^2 + 2m + 5)^2.$$

What is the order of the differential equation and find its general solution.

(10)

- (c) A particle rests in equilibrium under the reaction of two centers of forces which attract directly as the distance, their intensity being  $\mu, \mu'$ , the particle is displaced slightly towards one of them, show that the time of a small oscillation is

$$T = \frac{2\pi}{\sqrt{\mu + \mu'}}$$

(10)

- (d) A solid sphere rests inside a fixed rough hemispherical bowl of twice its radius. Show that however large a weight is attached to the highest point of the sphere, the equilibrium is stable.

(10)

- (e) Find an equation for the plane passing through the points  $P_1(3, 1, -2)$ ,  $P_2(-1, 2, 4)$ ,  $P_3(2, -1, 1)$  by using vector method.

(10)

6. (a) Solve  $x^2 \left( \frac{dy}{dx} \right)^2 + y(2x + y) \frac{dy}{dx} + y^2 = 0$ .

(10)

- (b) Using differential equations show that the system of confocal conics given by

$$\frac{x^2}{a^2 + \lambda} + \frac{y^2}{b^2 + \lambda} = 1, \lambda \text{ real is}$$

Self-orthogonal.

(10)

(c) Solve

$$(1-x^2) \frac{d^2 y}{dx^2} - x \frac{dy}{dx} - a^2 y = 0$$

given that  $y = e^{\sin^{-1} x}$  is one solution of this equation. (10)

(d) Find a general solution of  $y'' + y = \tan x$ ,  $-\frac{\pi}{2} < x < \frac{\pi}{2}$  by variation of parameters. (10)

7. (a) A particle moves with central acceleration  $(\mu u^2 + \lambda u^3)$  and the velocity of projection at distance R is V, show that the particle will ultimately go off to infinity if

$$V^2 > \frac{2\mu}{R} + \frac{\lambda}{R^2}. \quad (13)$$

(b) A smooth parabolic wire is fixed with its axis vertical and vertex downwards and in it is placed a uniform rod of length  $2l$  with its ends resting on wire. Show that, for equilibrium, the rod is either horizontal or makes with horizontal an angle  $\theta$  given by  $\cos^2 \theta = \frac{2a}{l}$ ,  $4a$  being the latus rectum of the parabola. (14)

(c) Prove that if volume  $v$  and  $V$  of two different substances balance in vacuum and volumes  $v'$ ,  $V'$  balance when weighed in a liquid, the densities of the substances and the liquid are as

$$\frac{v' - V'}{v} : \frac{v' - v'}{V} :: \left( \frac{v'}{V} - \frac{V'}{V} \right) \quad (13)$$

8. (a) Prove that

$$\nabla \times (\nabla \times \vec{A}) = -\nabla^2 \vec{A} + \nabla(\nabla \cdot \vec{A}) \quad (10)$$

(b) If

$$\nabla \cdot \vec{E}, \nabla \cdot \vec{H}, \nabla \times \vec{E} = -\frac{\partial \vec{H}}{\partial t}, \nabla \times \vec{H} = \frac{\partial \vec{E}}{\partial t}$$

Show that  $\vec{E}$  and  $\vec{H}$  satisfy

$$\nabla^2 \vec{u} = \frac{\partial^2 \vec{u}}{\partial t^2} \quad (10)$$

(c) Given the space curve  $x=t, y=t^2, z=\frac{2}{3}t^3$ . Find (i) the curvature  $\rho$  (ii) the torsion  $\tau$ . (10)

(d) If  $\vec{F} = (y^2 + z^2 - x^2)\vec{i} + (z^2 + x^2 - y^2)\vec{j} + (x^2 + y^2 - z^2)\vec{k}$ , evaluate  $\iint_S \text{curl } \vec{F} \cdot \vec{n} ds$ , taken over the portion of the surface,

$$x^2 + y^2 - z^2 - 2ax + az = 0 \text{ above the plane } z = 0 \text{ and verify Stoke's theorem.} \quad (10)$$

# MATHEMATICS

## PAPER - II

### SECTION A

1. Answer any four parts

(4 x 10 = 40)

(a) Write the elements of the symmetric group  $S_3$  of degree 3, prepare its multiplication table and find all normal subgroups of  $S_3$

(b) Change the order of integration in the integral

$$\int_0^{ka} \int_{a^2/ka}^{ay/a} dy \, dx$$

and evaluate it.

(c) Compute the Taylor series around  $z = 0$  and give the radius of convergence for  $\frac{z}{z-1}$

(d) By graphical method solve the linear programming problem

$$\text{Maximize } Z = 100 X_1 + 40 X_2$$

$$\text{subject to } 5 X_1 + 2 X_2 \leq 1000$$

$$3 X_1 + 2 X_2 \leq 900$$

$$X_1 + 2 X_2 \leq 500$$

$$X_1, X_2 \geq 0$$

(e) If

$$a_n = \log \left( 1 + \frac{1}{n} \right) + \log \left( 1 + \frac{2}{n} \right) + \dots + \log \left( 1 + \frac{n}{n} \right)$$

Find  $\lim_{n \rightarrow \infty} a_n$

2. (a) If every element of a group  $G$  is its own inverse, prove that the group  $G$  is abelian. Is the converse true? Justify your claim.

(14)

(b) Discuss the maxima and minima of  $x^3 y^2 (1-x-y)$ .

(13)

(c) State the Weierstrass M-test for uniform convergence of an infinite series of functions. Prove that the series

$$\sum_{n=1}^{\infty} \frac{x}{n^n (1 + nx^2)} \text{ with } 0 < 1$$

is uniformly convergent on  $(-\infty, \infty)$ .

(13)

3. (a) Define a field and prove that every finite integral domain is a field.

(3 + 10)

(b) Show that the function  $f(z) = \sqrt{xy}$  is not regular at the origin although the Cauchy-Riemann equations are satisfied.

(13)



- (c) By using the Residue Theorem evaluate the integral

$$\int_0^{2\pi} \frac{d\theta}{1 - 2a \sin \theta + a^2} \text{ where } 0 < a < 1.$$

(14)

4. (a) Define a unique factorization domain. Show that  $Z[\sqrt{-5}]$  is an integral domain which is not a unique factorization domain.

(3+10)

- (b) Using simplex method solve the linear programming problem:

$$\text{Maximize } Z = X_1 + X_2 + 3X_3$$

$$\text{subject to } 3X_1 + 2X_2 + X_3 \leq 3$$

$$2X_1 + X_2 + 2X_3 \leq 2$$

$$X_1, X_2, X_3 \geq 0.$$

(13)

- (c) A company has three plants A, B and C, three ware houses X, Y and Z. The number of units available at the plants is 60, 70, 80 and the demands at X, Y, Z are 50, 80, 80 respectively. The unit cost of the transportation is given in the following table:-

	X	Y	Z
A	8	7	3
B	3	8	9
C	11	3	4

Find the allocation so that the total transportation cost is minimum.

(14)

## SECTION B

5. Answer any four parts:-

(4 × 10 = 40)

- (a) Find the complete integral of the partial differential equation

$$x^2 p^2 + y^2 q^2 = z^2.$$

- (b) Find Lagrange's interpolation polynomial  $P_2(x)$  which satisfies.

$$f(0) = P_2(0) = 1$$

$$f(-1) = P_2(-1) = 2$$

$$f(1) = P_2(1) = 3.$$

Find  $f(0.5)$ .

- (c) Convert the decimal number  $(1479.25)_{10}$  to the binary and the hexadecimal numbers.

- (d) Find the moment of inertia of the area bounded by  $r^2 = a^2 \cos 2\theta$  about its axis.

- (e) Show that  $\frac{x^2}{a^2} f(t) + \frac{y^2}{b^2} \cdot \frac{1}{f(t)} = 1$  is a possible form of the bounding surface of a liquid.

6. (a) Solve by Charpit's method

$$(p^2 + q^2) y = qz.$$

(13)

- (b) If  $\varphi(x)$  is a continuous and bounded function of  $-\infty < x < \infty$ , prove that the function  $u(x,t) = \frac{1}{2\sqrt{\pi kt}} \int_{-\infty}^{\infty} \varphi(\xi) e^{-(x-\xi)^2/4kt} d\xi$  is a solution of the initial value problem:

$$u_t = k u_{xx} = 0, \quad -\infty < x < \infty, t > 0$$

$$u(x,0) = \varphi(x) \text{ for } -\infty < x < \infty.$$

- (c) Two equal masses  $m_1$  and  $m_2$  with  $m_1 > m_2$  are suspended by a light string over a pulley of mass  $M$  and radius  $a$ . There is no slipping and the friction of axle may be neglected. If  $f$  be the acceleration, show that this is constant and if  $k^2$  be the radius of gyration of the pulley about the axis, show that

$$k^2 = \frac{a^2}{Mf} [(g-f)m_1 - (g+f)m_2]$$

(13)

7. (a) Four equal rods, each of length  $2a$ , are hinged at their ends so as to form a rhombus ABCD. The angles B and D are connected by an elastic string and the lowest end A rests on a horizontal plane whilst the end C slides on a smooth vertical wire passing through A. In the position of equilibrium the string is stretched to twice its natural length and the angle BAD is  $2\alpha$ . Show that the time of small oscillation about this position is

$$2\pi \left\{ \frac{2a(1+3\sin^2\alpha)\cos\alpha}{(3g\cos 2\alpha)} \right\}^{\frac{1}{2}}$$

(14)

- (b) If  $q$  is the resultant velocity at any point of a fluid which is moving irrotationally in two dimensions, prove that

$$\left( \frac{\partial q}{\partial x} \right)^2 + \left( \frac{\partial q}{\partial y} \right)^2 = q^2 \nabla^2 q$$

- (c) By applying the Newton-Raphson method to  $f(x) = x^2 - a$  where  $a > 0$ , prove that

$$x_{n+1} = \frac{1}{2} \left( x_n + \frac{a}{x_n} \right)$$

For  $n = 0, 1, 2, 3$  Apply this formula to find  $\sqrt{2}$

8. (a) Write an algorithm for generating even integers  $\leq 100$ . Also draw the flow chart which executes this algorithm.

(13)

- (b) Applying Simpson's one-third rule compute the value of the definite integral

$$\int_1^{1.2} \log x \, dx$$

with  $h = 0.2$  and estimate the error.

(13)

- (c) State the conditions under which the equations of motion can be integrated. Obtain Bernoulli's equation for the steady irrotational motion of an incompressible liquid.

(14)