

**MATHEMATICS**  
**Paper II**  
**(CONVENTIONAL)**

Time allowed : Three Hours

Maximum Marks : 200

**Question Paper Specific Instructions**

*Please read each of the following instructions carefully before attempting questions :*

*There are **EIGHT** questions in all, out of which **FIVE** are to be attempted.*

*Questions no. 1 and 5 are compulsory. Out of the remaining **SIX** questions, **THREE** are to be attempted selecting at least **ONE** question from each of the two Sections **A** and **B**.*

*Attempts of questions shall be counted in chronological order. Unless struck off, attempt of a question shall be counted even if attempted partly. Any page or portion of the page left blank in the Answer Book must be clearly struck off.*

*All questions carry equal marks. The number of marks carried by a question / part is indicated against it.*

*Answers must be written in **ENGLISH** only.*

*Unless otherwise mentioned, symbols and notations have their usual standard meanings.*

*Assume suitable data, if necessary and indicated the same clearly.*

**SECTION A**

**Q.1.** (a) Evaluate :

$$\lim_{x \rightarrow 0} \left( \frac{e^{ax} - e^{bx} + \tan x}{x} \right)$$

10

(b) Prove that if every element of a group  $(G, 0)$  be its own inverse, then it is an abelian group.

10

(c) Construct an analytic function

$$f(z) = u(x, y) + iv(x, y), \text{ where}$$

$$v(x, y) = 6xy - 5x + 3.$$

Express the result as a function of  $z$ .

10

(d) Find the optimal assignment cost from the following cost matrix :

10

	A	B	C	D
I	4	5	4	3
II	3	2	2	6
III	4	5	3	5
IV	2	4	2	6

**IMS-Institute of Mathematical Sciences**

- Q.2.** (a) Show that any finite integral domain is a field. 13  
 (b) Every field is an integral domain — Prove it. 13  
 (c) Solve the following Salesman problem : 14

	A	B	C	D
A	∞	12	10	15
B	16	∞	11	13
C	17	18	∞	20
D	13	11	18	∞

- Q.3.** (a) Show that the function  $f(x) = x^2$  is uniformly continuous in  $(0, 1)$  but not in  $\mathbb{R}$ . 13  
 (b) Prove that :  
 (i) the intersection of two ideals is an ideal.  
 (ii) a field has no proper ideals. 14

- (c) Evaluate  $\oint_c \frac{e^{2z}}{(z+1)^4} dz$  where  $c$  is the circle  $|z| = 3$ . 13

- Q.4.** (a) Find the area of the region between the x-axis and  $y = (x-1)^3$  from  $x = 0$  to  $x = 2$ . 13  
 (b) Find Laurent series about the indicated singularity. Name the singularity and give the region of convergence.

$\frac{z - \sin z}{z^3}; z = 0.$  13

- (c)  $x_1 = 4, x_2 = 1, x_3 = 3$  is a feasible solution of the system of equations  
 $2x_1 - 3x_2 + x_3 = 8$   
 $x_1 + 2x_2 + 3x_3 = 15$

Reduce the feasible solution to two different basic feasible solutions. 14

**IMS-Institute of Mathematical Sciences**

**SECTION B**

- Q.5.** (a) Use Newton – Raphson method and derive the iteration scheme  $x_{n+1} = \frac{1}{2} \left( x_n + \frac{N}{x_n} \right)$  to calculate an approximate value of the square root of a number N. Show that the formula  $\sqrt{N} \approx \frac{A+B}{4} + \frac{N}{A+B}$  where  $AB = N$ , can easily be obtained if the above scheme is applied two times. Assume  $A = 1$  as an initial guess value and use the formula twice to calculate the value of  $\sqrt{2}$  [For 2<sup>nd</sup> iteration, one may take  $A =$  result of the 1<sup>st</sup> iteration]. 14
- (b) Eliminate the arbitrary function f from the given equation 12  
 $f(x^2 + y^2 + z^2, x + y + z) = 0$
- (c) Derive the Hamiltonian and equation of motion for a simple pendulum. 14
- Q.6.** (a) Solve the PDE : 12  
 $xu_x + yu_y + zu_z = xyz$
- (b) Convert  $(0.231)_5$ ,  $(104.231)_5$  and  $(247)_7$  to base 10. 12
- (c) Rewrite the hyperbolic equation  $x^2u_{xx} - y^2u_{yy} = 0$  ( $x > 0, y > 0$ ) in canonical form. 16
- Q.7.** (a) Find the values of a and b in the 2-D velocity field  $\vec{v} = (3y^2 - ax^2)\hat{i} + bxy\hat{j}$  so that the flow becomes incompressible and irrotational. Find the stream function of the flow. 14
- (b) Write an algorithm to find the inverse of a given non-singular diagonally dominant square matrix using Gauss – Jordan method. 13
- (c) Find the solution of the equation 13  
 $\left(\frac{\partial u}{\partial x}\right)^2 + \left(\frac{\partial u}{\partial y}\right)^2 = 1$   
 that passes through the circle  
 $x^2 + y^2 = 1, u = 1.$
- Q.8.** (a) Solve the following heat equation, using the method of separation of variables : 16  
 $\frac{\partial u}{\partial t} = \frac{\partial^2 u}{\partial x^2}, 0 < x < 1, t > 0$   
 subject to the conditions  
 $u = 0$  at  $x = 0$  and  $x = 1$ , for  $t > 0$   
 $u = 4x(1 - x)$ , at  $t = 0$  for  $0 \leq x \leq 1.$
- (b) Use the Classical Fourth-order Runge – Kutta method with  $h = 0.2$  to calculate a solution at  $x = 0.4$  for the initial value problem  $\frac{du}{dx} = 4 - x^2 + u, u(0) = 0$  on the interval  $[0, 0.4]$ . 12
- (c) Draw a flow chart for testing whether a given real number is a prime or not. 12