




UPSC-CSE 2019

Mains

MATHEMATICS

Optional Paper-I

Solutions

SECTION - A

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

1.(a) → Let $f: [0, \pi/2] \rightarrow \mathbb{R}$ be a continuous function such that

$$f(x) = \frac{\cos^2 x}{4x^2 - \pi^2}, \quad 0 \leq x < \pi/2$$

Find the value of $f(\pi/2)$.

Solution :

Given that

$f: [0, \pi/2] \rightarrow \mathbb{R}$ is a continuous function such that $f(x) = \frac{\cos^2 x}{4x^2 - \pi^2}$,

$$0 \leq x < \pi/2.$$

Now, since f is continuous on $[0, \pi/2]$

$$\text{We have } \lim_{x \rightarrow \pi/2^-} f(x) = f(\pi/2)$$

$$\begin{aligned} \Rightarrow f(\pi/2) &= \lim_{x \rightarrow \pi/2^-} \frac{\cos^2 x}{4x^2 - \pi^2} \quad \left(\frac{0}{0}\right) \\ &= \lim_{x \rightarrow \pi/2^-} \frac{2 \cos x (-\sin x)}{8x} \end{aligned}$$

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

$$= \lim_{x \rightarrow \pi/2} - \frac{\sin 2x}{8x}$$

$$= \frac{-\sin 2(\pi/2)}{8(\pi/2)}$$

$$= 0$$

$\therefore$ The value of $f(\pi/2) = 0$.

1.(b) → Let $f: D(\subseteq \mathbb{R}^2) \rightarrow \mathbb{R}$ be a function and $(a, b) \in D$. If $f(x, y)$ is continuous at (a, b) , then show that the functions $f(x, b)$ and $f(a, y)$ are continuous at $x=a$ and at $y=b$ respectively.

Solution:

Let $f: D(\subseteq \mathbb{R}^2) \rightarrow \mathbb{R}$ be a function and $(a, b) \in D$.

If $f(x, y)$ is continuous at (a, b) then by definition,

$$\lim_{(x, y) \rightarrow (a, b)} f(x, y) = f(a, b)$$

i.e. $f(x, y) \rightarrow f(a, b)$ as $(x, y) \rightarrow (a, b)$

i.e. for given $\epsilon > 0$ (however small)

$\exists \delta > 0$ ($\delta(\epsilon)$) such that

$$|f(x, y) - f(a, b)| < \epsilon \text{ whenever } \|(x, y) - (a, b)\| < \delta.$$

A point to be particularly noticed is that if a function of more than

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

one variable is continuous at a point, it is continuous at that point when considered as a function of single variable.

To be more specific if a function f of two variables x, y is continuous at (a, b) then $f(x, b)$ is continuous at $x=a$ and $f(a, y)$ that of f at $y=b$.

Hence, the result.

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Ques: 1(c) Let $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear map such that $T(2,1) = (5,7)$ and $T(1,2) = (3,3)$. If A is the matrix corresponding of T with respect to the standard bases e_1, e_2 , then find Rank (A) .

Solution:-

Let $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear map such that

$$T(2,1) = (5,7) \quad \text{and} \quad T(1,2) = (3,3).$$

Let us find the matrix A corresponding to T with respect to the standard bases e_1, e_2 .

where $e_1 = (1,0)$
and $e_2 = (0,1)$.

Then;

$$T(2,1) = (5,7) = 5(1,0) + 7(0,1)$$

$$T(1,2) = (3,3) = 3(1,0) + 3(0,1)$$

$$\therefore A = \begin{bmatrix} 5 & 3 \\ 7 & 3 \end{bmatrix} \text{ is a matrix of } T.$$

$$\text{Since } |A| = \begin{vmatrix} 5 & 3 \\ 7 & 3 \end{vmatrix} = 15 - 21 = -6 \neq 0$$

$$\therefore \text{Rank of } A = 2.$$

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Ques: 1) d) If $A = \begin{bmatrix} 1 & 2 & 1 \\ 1 & -4 & 1 \\ 3 & 0 & -3 \end{bmatrix}$ and $B = \begin{bmatrix} 2 & 1 & 1 \\ 1 & -1 & 0 \\ 2 & 1 & -1 \end{bmatrix}$

then show that $AB = 6I_3$. Use this result to solve the following system of equations:

$$\begin{aligned} 2x + y + z &= 5 \\ x - y &= 0 \\ 2x + y - z &= 1. \end{aligned}$$

Solution:-

Given; $A = \begin{bmatrix} 1 & 2 & 1 \\ 1 & -4 & 1 \\ 3 & 0 & -3 \end{bmatrix}$ and $B = \begin{bmatrix} 2 & 1 & 1 \\ 1 & -1 & 0 \\ 2 & 1 & -1 \end{bmatrix}$

$$A \cdot B = \begin{bmatrix} 1 & 2 & 1 \\ 1 & -4 & 1 \\ 3 & 0 & -3 \end{bmatrix} \begin{bmatrix} 2 & 1 & 1 \\ 1 & -1 & 0 \\ 2 & 1 & -1 \end{bmatrix} = \begin{bmatrix} 6 & 0 & 0 \\ 0 & 6 & 0 \\ 0 & 0 & 6 \end{bmatrix}$$

$$A \cdot B = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = 6I_3$$

Hence proved.

$$A \cdot B = 6I_3$$

$$A \cdot B \cdot B^{-1} = 6I_3 B^{-1}$$

$$A = 6B^{-1}$$

$$B^{-1} = \frac{1}{6} A$$

$$2x + y + z = 5$$

$$x - y = 0$$

$$2x + y - z = 1$$

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

$$B = \begin{bmatrix} 2 & 1 & 1 \\ 1 & -1 & 0 \\ 2 & 1 & -1 \end{bmatrix} ; C = \begin{bmatrix} 5 \\ 0 \\ 1 \end{bmatrix} ; X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$$

and from the linear equation

$$BX = C \Rightarrow B^{-1}BX = B^{-1}C.$$

$$X = B^{-1}C$$

$$X = \frac{1}{6}AC$$

$$X = \frac{1}{6} \begin{bmatrix} 1 & 2 & 1 \\ 1 & -4 & 1 \\ 3 & 0 & -3 \end{bmatrix} \begin{bmatrix} 5 \\ 0 \\ 1 \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \frac{1}{6} \begin{bmatrix} 6 \\ 6 \\ 12 \end{bmatrix} = \frac{1}{6} \times 6 \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix}$$

$$\therefore \underline{x=1, y=1, z=2.}$$

is the required result.

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

1.(e)

→ Show that the lines

$$\frac{x+1}{-3} = \frac{y-3}{2} = \frac{z+2}{1} \quad \text{and} \quad \frac{x}{1} = \frac{y-7}{-3} = \frac{z+7}{2}$$

intersect. Find the coordinates of the point of intersection and the equation of the plane containing them.

Solution: Part (a):

Any point on the line

$$\frac{x+1}{-3} = \frac{y-3}{2} = \frac{z+2}{1} \quad \text{is}$$

$$(-1-3r, 3+2r, -2+r) \quad \text{--- (1)}$$

Similarly, any point on the line

$$\frac{x}{1} = \frac{y-7}{-3} = \frac{z+7}{2} \quad \text{is}$$

$$(r', 7-3r', -7+2r') \quad \text{--- (2)}$$

If the two given lines intersect then for some value of r and r' the two

above points (i) and (ii) must coincide

$$\begin{aligned}\text{i.e. } -1-3r &= r'; \\ 3+2r &= 7-3r'; \\ -2+r &= -7+2r'\end{aligned}$$

Solving the first two of these equations, we get $r = -1$, $r' = 2$.

These values of r and r' satisfy the third equation also, hence the given lines intersect.

Substituting these values r and r' in (1) or (2) we get the required coordinates of the point of intersection as $(2, 1, -3)$. [I]

Part (b):

Also the equation of the plane containing the given lines is

$$\begin{vmatrix} x+1 & y-3 & z+2 \\ -3 & 2 & 1 \\ 1 & -3 & 2 \end{vmatrix} = 0$$

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

$$\Rightarrow (x+1)(4+3) - (y-3)(-6-1) + (z+2)(9-2) \\ = 0$$

$$\Rightarrow \boxed{x+y+z=0}$$

which is the required equation.

— [1]

2.(b) $\rightarrow$ Let A and B be two orthogonal matrices of same order and $\det A + \det B = 0$. Show that $A+B$ is a singular matrix.

Solution :

Given that A and B are two orthogonal matrices of same order and $\det A + \det B = 0$. We know that, all orthogonal matrices are invertible, so we can write

$$A+B = A(I+A^{-1}B) \quad \text{--- (1)}$$

Now, it is enough to show that $(I+A^{-1}B)$ is singular.

[$\because$ the product of matrices is singular as long as at least one of the matrices is singular.]

$$\begin{aligned} \text{Also, } \det(A^{-1} * B) &= \det(A^{-1}) * \det B \\ &= \det(A^{-1}) * (-\det(A)) \end{aligned}$$

$$[\because \det(A) + \det(B) = 0]$$

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Now,
since the determinant of an orthogonal matrix
is either 1 or -1,

$$\det(A)^{-1} * (-\det(A)) = -1.$$

[$\because$ This product is the product
of -1 and 1 (in some
order)]

————— (2)

Using (2), we have

$$\begin{aligned}\det(I + A^{-1}B) &= \det I + \det(A^{-1}B) \\ &= \det I + (-1) \\ &= 1 - 1 \\ &= 0\end{aligned}$$

$\Rightarrow A+B$ is singular matrix

[$\because$ A matrix is singular
 $\Leftrightarrow$ its determinant is 0].

Hence, proved.

2.(c) →

- (i) The plane $x+2y+3z=12$ cuts the axes of co-ordinates in A, B, C. Find the equations of the circle circumscribing the triangle ABC.

Solutions:

The given plane $x+2y+3z=12$ ——— (1)
meets the x-axis i.e., $y=0, z=0$ in the point A whose coordinates are $(12, 0, 0)$.

Similarly the co-ordinates of B and C where the given plane meets y-axis are $(0, 6, 0)$ and z-axis are $(0, 0, 4)$.

Thus, the point A, B, C are $(12, 0, 0)$, $(0, 6, 0)$ and $(0, 0, 4)$ respectively.

Let the equation of the circle circumscribing the triangle ABC be

$$x^2 + y^2 + z^2 + 2ux + 2vy + 2wz + d = 0$$

———— (2)

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

If it passes through A, B, C then we have

$$(12)^2 + 2u(12) + d = 0 \Rightarrow 24u + d + 144 = 0 \quad \text{--- (3)}$$

$$(6)^2 + 2v(6) + d = 0 \Rightarrow 12v + d + 36 = 0 \quad \text{--- (4)}$$

$$\text{and } (4)^2 + 2w(4) + d = 0 \Rightarrow 8w + d + 16 = 0 \quad \text{--- (5)}$$

From (3), (4) and (5) we get

$$2u = -[12 - (d/12)],$$

$$2v = -[6 - (d/6)] \quad \text{and}$$

$$2w = -[4 - (d/4)].$$

Substituting these values of $2u$, $2v$, $2w$ in (2), we get the required equation as

$$\underline{x^2 + y^2 + z^2 - [12 - (d/12)]x - [6 - (d/6)]y - [4 - (d/4)]z + d = 0}$$

where d can take any value.

Hence, the result.

2. (c) →
 (ii)

Prove that the plane $z=0$ cuts the enveloping cone of the sphere $x^2+y^2+z^2=11$ which has the vertex at $(2, 4, 1)$ in a rectangular hyperbola.

Solution:

The equation of the sphere is

$$x^2 + y^2 + z^2 - 11 = 0 \quad \text{and the vertex is } (2, 4, 1).$$

Here,

$$S = x^2 + y^2 + z^2 - 11, \quad x_1 = 2, y_1 = 4, z_1 = 1$$

$$S_1 = x_1^2 + y_1^2 + z_1^2 - 11 = 4 + 16 + 1 - 11$$

$$\text{i.e. } S_1 = x_1^2 + y_1^2 + z_1^2 - 11 = 10$$

$$\text{and } T = xx_1 + yy_1 + zz_1 - 11$$

$$= 2x + 4y + z - 11.$$

∴ Equation of the enveloping cone is

$$(x^2 + y^2 + z^2 - 11)(10) = (2x + 4y + z - 11)^2$$

[Using $SS_1 = T^2$]

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

$$\Rightarrow 10(x^2 + y^2 + z^2 - 11) - (2x + 4y + z - 11)^2 = 0$$

This meets the plane $z=0$ in the curve

$$10(x^2 + y^2 - 11) - (2x + 4y - 11)^2 = 0$$

$$\Rightarrow 10(x^2 + y^2) - (4x^2 + 16y^2 + \dots) = 0$$

This represents a rectangular hyperbola
in the XY -plane if co-eff. of $x^2 +$

$$\text{co. eff. of } y^2 = 0$$

$$\Rightarrow \text{if } (10 - 4) + (10 - 16) = 0$$

which is true

Hence, the result.

3.(a)

→ Find the maximum and the minimum value of the function $f(x) = 2x^3 - 9x^2 + 12x + 6$ on the interval $[2, 3]$.

Solution:

$$\text{Given } f(x) = 2x^3 - 9x^2 + 12x + 6$$

$$f'(x) = 6x^2 - 18x + 12$$

$$\text{Let } f'(x) = 0$$

$$6x^2 - 18x + 12 = 0$$

$$\Rightarrow x^2 - 3x + 2 = 0$$

$$\Rightarrow (x-1)(x-2) = 0$$

$$\Rightarrow x = 1, 2.$$

These are the critical points of the $f(x)$ which $\notin [2, 3]$. Thus, $f(x)$ is monotonic on $[2, 3]$.

Now, to check whether it is monotonic increasing or decreasing -

$$f'(2) = 6(4) - 18(2) + 12 = 24 - 36 + 12 = 0$$

$$f'(3) = 6(9) - 18(3) + 12 = 54 - 54 + 12 = 12 > 0$$

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Since, $f'(2) = 0$, $f'(3) > 0$ and

$$f'(3) > f'(2).$$

Hence, $f(x)$ is monotonically increasing on $[2, 3]$.

$$\begin{aligned} \text{Thus, } f_{\min} = f(x) \Big|_{x=2} &= 2 \times (2)^3 - 9 \times 4 + 12 \times 2 + 6 \\ &= 16 - 36 + 24 + 6 \\ &= 10. \end{aligned}$$

$$\begin{aligned} f_{\max} = f(x) \Big|_{x=3} &= 2 \times (3)^3 - 9 \times 9 + 12 \times 3 + 6 \\ &= 54 - 81 + 36 + 6 \\ &= 96 - 81 \\ &= 15. \end{aligned}$$

$$\therefore f_{\min} = 10 \text{ and } f_{\max} = 15.$$

Hence, the result.

3.(b) → Prove that, in general, three normals can be drawn from a given point to the paraboloid $x^2 + y^2 = 2az$, but if the point lies on the surface

$27a(x^2 + y^2) + 8(a - z)^3 = 0$ then two of the three normals coincide.

Solution:

The equations of the normal at (x_1, y_1, z_1) to the paraboloid $x^2 + y^2 = 2az$ are

$$\frac{x - x_1}{x_1} = \frac{y - y_1}{y_1} = \frac{z - z_1}{-a}$$

This passes through a given point (α, β, γ) if

$$\frac{\alpha - x_1}{x_1} = \frac{\beta - y_1}{y_1} = \frac{\gamma - z_1}{-a} = \lambda \text{ (say)}$$

$$\left. \begin{aligned} \text{This gives } \alpha - x_1 &= \lambda x_1 \Rightarrow x_1 = \alpha / (1 + \lambda) \\ \text{Similarly, } y_1 &= \beta / (1 + \lambda), \quad z_1 = \gamma + a\lambda \end{aligned} \right\} \quad (1)$$

Also (x_1, y_1, z_1) lies on the given paraboloid, so

$$x_1^2 + y_1^2 = 2az_1 \Rightarrow \left[\frac{\alpha}{1 + \lambda} \right]^2 + \left[\frac{\beta}{1 + \lambda} \right]^2 = 2a(\gamma + a\lambda),$$

[from (1)]

$$\Rightarrow \alpha^2 + \beta^2 = 2a(\gamma + a\lambda)(1 + \lambda)^2 \quad \text{--- (2)}$$

This being a cubic in λ gives three values of λ and so from (1) there are three points on the paraboloid normals at which pass through (α, β, γ) .

The equation (2) can be rewritten as

$$f(\lambda) \equiv 2a(1 + \lambda)^2(\gamma + a\lambda) - (\alpha^2 + \beta^2) = 0 \quad \text{--- (3)}$$

The condition that this equation has two equal roots is obtained by eliminating λ between $f(\lambda) = 0$ and $f'(\lambda) = 0$.

$$\text{From (3), } f'(\lambda) = 0 \text{ means } 2a(1 + \lambda)^2(a) + 4a(1 + \lambda)(\gamma + a\lambda) = 0$$

$$\Rightarrow a(1 + \lambda) + 2(\gamma + a\lambda) = 0 \quad (\because 1 + \lambda \neq 0)$$

$$\Rightarrow (a + 2\gamma) + \lambda(3a) = 0$$

$$\Rightarrow \lambda = -(a + 2\gamma) / (3a)$$

Substituting this value of λ in (3), we get,

$$2a \left[1 - \frac{a + 2\gamma}{3a} \right]^2 \left[\gamma - \frac{a(a + 2\gamma)}{3a} \right] = \alpha^2 + \beta^2$$

$$\Rightarrow 2a [2(a-r)]^2 [a(r-a)] = 27a^3 (\alpha^2 + \beta^2)$$

$$\Rightarrow 27a(\alpha^2 + \beta^2) + 8(a-r)^3 = 0$$

$\therefore$ Locus of the point (α, β, r) is

$$27a(x^2 + y^2) + 8(a-z)^3 = 0.$$

Hence, proved.

3.(c)

Let

$$A = \begin{bmatrix} 5 & 7 & 2 & 1 \\ 1 & 1 & -8 & 1 \\ 2 & 3 & 5 & 0 \\ 3 & 4 & -3 & 1 \end{bmatrix}$$

(i) Find the rank of matrix A.

(ii) Find the dimension of the subspace

$$V = \left\{ (x_1, x_2, x_3, x_4) \in \mathbb{R}^4 \mid A \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = 0 \right\}.$$

Solution:

The given matrix is

$$A = \begin{bmatrix} 5 & 7 & 2 & 1 \\ 1 & 1 & -8 & 1 \\ 2 & 3 & 5 & 0 \\ 3 & 4 & -3 & 1 \end{bmatrix}$$

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Now, we use row transformations to obtain an echelon matrix.

$$\sim \begin{bmatrix} 1 & 1 & -8 & 1 \\ 2 & 3 & 5 & 0 \\ 3 & 4 & -3 & 1 \\ 5 & 7 & 2 & 1 \end{bmatrix}$$

$$R_1 \leftrightarrow R_2 ;$$

$$R_2 \leftrightarrow R_3 ;$$

$$R_3 \leftrightarrow R_4 .$$

$$\sim \begin{bmatrix} 1 & 1 & -8 & 1 \\ 0 & 1 & 21 & -2 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$$R_2 \rightarrow R_2 - 2R_1 .$$

$$R_3 \rightarrow R_3 - (R_1 + R_2)$$

$$R_4 \rightarrow R_4 - (R_2 + R_3) \quad \text{--- (I)}$$

which is clearly in the echelon form and the no. of non-zero is 2.

$$\therefore \rho(A) = 2.$$

--- (V)

Let $\mathbb{R}^4 = \{(x_1, x_2, x_3, x_4) / x_1, x_2, x_3, x_4 \in \mathbb{R}\}$ be given vector space.

$$\text{Let } V = \left\{ (x_1, x_2, x_3, x_4) \in \mathbb{R}^4 / A \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = 0 \right\} \subseteq \mathbb{R}^4$$

We have $AX = 0$

$$\Rightarrow \begin{bmatrix} 5 & 7 & 2 & 1 \\ 1 & 1 & -8 & 1 \\ 2 & 3 & 5 & 0 \\ 3 & 4 & -3 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

$$\Rightarrow \begin{bmatrix} 1 & 1 & -8 & 1 \\ 0 & 1 & 21 & -2 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix} \quad [\text{from (I)}]$$

$$\Rightarrow x_1 + x_2 - 8x_3 + x_4 = 0 \quad \text{--- (1)}$$

$$x_2 + 21x_3 - 2x_4 = 0 \quad \text{--- (2)}$$

from (2),

$$x_2 = -21x_3 + 2x_4$$

$\therefore$ from (1)

$$x_1 - 21x_3 + 2x_4 - 8x_3 + x_4 = 0$$

$$\Rightarrow x_1 - 29x_3 + 3x_4 = 0$$

$$\Rightarrow x_1 = 29x_3 - 3x_4$$

$\therefore$ There are two free variables x_3 and x_4 (say).

$\therefore$ $\dim V = 2$ and

$$V = \left\{ (29x_3 - 3x_4, -21x_3 + 2x_4, x_3, x_4) / \right. \\ \left. x_3, x_4 \in \mathbb{R} \right\} \quad \text{--- (ii)}$$

Hence, the result.

4.(a) → State the Cayley-Hamilton theorem.
Use this theorem to find A^{100} , where

$$A = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$$

Solution:

Statement:

“Cayley-Hamilton Theorem”

Every square matrix A is a zero of its characteristic polynomial.

The given matrix is

$$A = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$$

The characteristic polynomial for the given matrix $A = |A - \lambda I|$

$$= (1-\lambda)(\lambda^2-1)$$

$$= -\lambda^3 + \lambda^2 + \lambda - 1$$

By Cayley-Hamilton Theorem, we have

$$-A^3 + A^2 + A - I = 0$$

$$\Rightarrow A^3 = A^2 + A - I \quad \text{--- (1)}$$

Now,

$$A^4 = A^3 + A^2 - A$$

$$= (A^2 + A - I) + A^2 - A \quad \text{--- (using (1))}$$

$$A^4 = 2A^2 - I$$

Similarly,

$$A^5 = 2A^2 + A - 2I ;$$

$$A^6 = 3A^2 + 3A - 5I \quad \text{--- (2)}$$

$\vdots$

$$A^8 = 4A^2 + 3A - 6I \quad \text{--- (3)}$$

$\vdots$

$$A^{10} = 5A^2 + 3A - 7I. \quad \text{--- (4)}$$

$\vdots$

By inductive sequence, we have

$$A^{2n} = nA^2 + 3A - (n+2)I. \quad \text{--- (.)}$$

$$\therefore A^{100} = 50A^2 + 3A - 52I.$$

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

$$\therefore A^{100} = 50 \begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix} + 3 \begin{bmatrix} 1 & 0 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} - 52 \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} 53 & 0 & 0 \\ 53 & 50 & 3 \\ 50 & 3 & 50 \end{bmatrix} - \begin{bmatrix} 52 & 0 & 0 \\ 0 & 52 & 0 \\ 0 & 0 & 52 \end{bmatrix}$$

$$\therefore A^{100} = \begin{bmatrix} 1 & 0 & 0 \\ 53 & -2 & 3 \\ 50 & 3 & -2 \end{bmatrix}$$

which is the required matrix.

Hence, the result.

4.(b) → Find the length of the normal chord through a point P of the ellipsoid

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1 \quad \text{and prove that}$$

if it is equal to $4PG_3$, where G_3 is the point where the normal chord through P meets the xy -plane, then P lies on the cone

$$\frac{x^2}{a^6} (2c^2 - a^2) + \frac{y^2}{b^6} (2c^2 - b^2) + \frac{z^2}{c^4} = 0.$$

Solution:

Let P be (α, β, γ) , then the equations of the normal to the given ellipsoid at $P(\alpha, \beta, \gamma)$

are $\frac{x-\alpha}{(p\alpha/a^2)} = \frac{y-\beta}{(p\beta/b^2)} = \frac{z-\gamma}{(p\gamma/c^2)} = r(\text{say})$ — (1)

where $\frac{1}{p^2} = \frac{\alpha^2}{a^4} + \frac{\beta^2}{b^4} + \frac{\gamma^2}{c^4}$ — (2)

∴ The co-ordinates of any point Q on the normal (1) are

$$\left(\alpha + \frac{p\alpha}{a^2} r, \beta + \frac{p\beta}{b^2} r, \gamma + \frac{p\gamma}{c^2} r \right),$$

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

where r is the distance of Q from P .

If Q lies on the given ellipsoid i.e. PQ is the normal chord, then

$$\frac{1}{a^2} \left(\alpha + \frac{p\alpha}{a^2} r \right)^2 + \frac{1}{b^2} \left(\beta + \frac{p\beta}{b^2} r \right)^2 + \frac{1}{c^2} \left(\gamma + \frac{p\gamma}{c^2} r \right)^2 = 1.$$

$$\Rightarrow r^2 p^2 \left(\frac{\alpha^2}{a^6} + \frac{\beta^2}{b^6} + \frac{\gamma^2}{c^6} \right) + 2rp \left(\frac{\alpha^2}{a^4} + \frac{\beta^2}{b^4} + \frac{\gamma^2}{c^4} \right) +$$

$$\left(\frac{\alpha^2}{a^2} + \frac{\beta^2}{b^2} + \frac{\gamma^2}{c^2} \right) = 1$$

$$\Rightarrow r^2 p^2 \left(\frac{\alpha^2}{a^6} + \frac{\beta^2}{b^6} + \frac{\gamma^2}{c^6} \right) + 2rp \left(\frac{1}{p^2} \right) = 0$$

[from (2) and
 $\sum \frac{\alpha^2}{a^2} = 1$

as $P(\alpha, \beta, \gamma)$ lies on
 the given conicoid.]

$$\Rightarrow \boxed{r = \frac{-2}{p^3 \left(\frac{\alpha^2}{a^6} + \frac{\beta^2}{b^6} + \frac{\gamma^2}{c^6} \right)}} = \text{length of normal chord } PQ$$

— (3)

Also,

Let the normal at $P(\alpha, \beta, \gamma)$ meets the coordinate planes viz. yz , zx and xy planes at G_1 , G_2 and G_3 , then putting $x=0$, $y=0$ and $z=0$ in succession in the Eqⁿ (1), we have respectively,

$$PG_1 = -a^2/p, \quad PG_2 = -b^2/p \quad \text{and} \quad PG_3 = -c^2/p \quad \text{--- (4)}$$

Given $PQ = 4PG_3$

$$\Rightarrow PQ = 4(-c^2/p)$$

$$\Rightarrow \frac{-2}{p^3 \left(\frac{\alpha^2}{a^6} + \frac{\beta^2}{b^6} + \frac{\gamma^2}{c^6} \right)} = 4 \left(-\frac{c^2}{p} \right)$$

$$\Rightarrow 2c^2 \left(\frac{\alpha^2}{a^6} + \frac{\beta^2}{b^6} + \frac{\gamma^2}{c^6} \right) = \frac{1}{p^2} = \frac{\alpha^2}{a^4} + \frac{\beta^2}{b^4} + \frac{\gamma^2}{c^4} \quad \text{--- from (2)}$$

$$\Rightarrow \frac{\alpha^2}{a^6} (2c^2 - a^2) + \frac{\beta^2}{b^6} (2c^2 - b^2) + \frac{\gamma^2}{c^6} (2c^2 - c^2) = 0$$

$\therefore$ The locus of $P(\alpha, \beta, \gamma)$ is

$$\frac{x^2}{a^6} (2c^2 - a^2) + \frac{y^2}{b^6} (2c^2 - b^2) + \frac{z^2}{c^4} = 0$$

Hence, proved.

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

4. (c) →

(i) If $u = \sin^{-1} \sqrt{\frac{x^{1/3} + y^{1/3}}{x^{1/2} + y^{1/2}}}$ then

show that $\sin^2 u$ is a homogeneous function of x and y of degree $-1/6$.
Hence show that

$$x^2 \frac{\partial^2 u}{\partial x^2} + 2xy \frac{\partial^2 u}{\partial x \partial y} + y^2 \frac{\partial^2 u}{\partial y^2} = \frac{\tan u}{12} \left(\frac{13}{12} + \frac{\tan^2 u}{12} \right)$$

Solution:

Given that $u = \sin^{-1} \sqrt{\frac{x^{1/3} + y^{1/3}}{x^{1/2} + y^{1/2}}}$, we can

write

$$\sin u = \left[\frac{x^{1/3} + y^{1/3}}{x^{1/2} + y^{1/2}} \right]^{1/2} = \frac{x^{1/6}}{x^{1/4}} \left[\frac{1 + (y/x)^{1/3}}{1 + (y/x)^{1/2}} \right]^{1/2}$$

$$= x^{-1/12} f(y/x)$$

$$\Rightarrow \sin^2 u = \left[\frac{x^{1/3} + y^{1/3}}{x^{1/2} + y^{1/2}} \right] = \frac{x^{1/3}}{x^{1/2}} \left[\frac{1 + (y/x)^{1/3}}{1 + (y/x)^{1/2}} \right]$$

$$= x^{-1/6} f(y/x)$$

Thus, $z = \sin^2 u$ is a homogeneous function of x and y of degree $-1/6$. — (i)

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Now, by Euler's theorem,

$$x \frac{\partial z}{\partial x} + y \frac{\partial z}{\partial y} = -\frac{1}{12} z, \text{ where } z = \sin u$$

$$\Rightarrow x \cos u \frac{\partial u}{\partial x} + y \cos u \frac{\partial u}{\partial y} = -\frac{1}{12} \sin u$$

$$\Rightarrow x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} = -\frac{1}{12} \tan u \quad \text{--- (1)}$$

Differentiating (1) partially w.r.t. x and y , respectively

$$x \frac{\partial^2 u}{\partial x^2} + \frac{\partial u}{\partial x} + y \frac{\partial^2 u}{\partial x \partial y} = -\frac{1}{12} \sec^2 u \frac{\partial u}{\partial x} \quad \text{--- (2)}$$

$$x \frac{\partial^2 u}{\partial x \partial y} + y \frac{\partial^2 u}{\partial y^2} + \frac{\partial u}{\partial y} = -\frac{1}{12} \sec^2 u \frac{\partial u}{\partial y} \quad \text{--- (3)}$$

Multiply (2) by x , (3) by y and add to get

$$\left(x^2 \frac{\partial^2 u}{\partial x^2} + 2xy \frac{\partial^2 u}{\partial x \partial y} + y^2 \frac{\partial^2 u}{\partial y^2} \right) + \left(x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} \right) = -\frac{1}{12} \sec^2 u \left(x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} \right) \quad \text{--- (4)}$$

From (1) and (4), we get

$$\begin{aligned} x^2 \frac{\partial^2 u}{\partial x^2} + 2xy \frac{\partial^2 u}{\partial x \partial y} + y^2 \frac{\partial^2 u}{\partial y^2} &= \frac{1}{12} \tan u \left(1 + \frac{1}{12} (\sec^2 u) \right) \\ &= \frac{\tan u}{12} \left(\frac{12 + 1 + \sec^2 u}{12} \right) \\ &= \frac{\tan u}{12} \left(\frac{13 + \tan^2 u}{12} \right) \end{aligned}$$

Hence, Proved.

4.(c)
(ii)

Using the Jacobian method, show that if $f'(x) = \frac{1}{1+x^2}$ and $f(0) = 0$, then

$$f(x) + f(y) = f\left(\frac{x+y}{1-xy}\right).$$

Solution:

Given that

$$f'(x) = \frac{1}{1+x^2} \quad \text{and} \quad f(0) = 0.$$

$$\text{Let } u = f(x) + f(y) \quad \text{and} \quad v = \frac{x+y}{1-xy}$$

— (1) — (2)

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

We have

$$\begin{aligned} \frac{\partial(u, v)}{\partial(x, y)} &= \begin{vmatrix} \frac{\partial u}{\partial x} & \frac{\partial u}{\partial y} \\ \frac{\partial v}{\partial x} & \frac{\partial v}{\partial y} \end{vmatrix} \\ &= \begin{vmatrix} f'(x) & f'(y) \\ \frac{(1-xy) - (x+y)(-y)}{(1-xy)^2} & \frac{(1-xy) - (x+y)(-x)}{(1-xy)^2} \end{vmatrix} \\ &= \begin{vmatrix} \frac{1}{1+x^2} & \frac{1}{1+y^2} \\ \frac{1+y^2}{(1-xy)^2} & \frac{1+x^2}{(1-xy)^2} \end{vmatrix} \\ &= \frac{1}{(1-xy)^2} - \frac{1}{(1-xy)^2} \\ &= 0 \end{aligned}$$

Since, $\frac{\partial(u, v)}{\partial(x, y)} = 0$.

$\therefore$ The given functions are not independent.
 i.e. the functions u and v are functionally related.

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Let $u = \phi(v)$ then

$$f(x) + f(y) = \phi\left(\frac{x+y}{1-xy}\right)$$

for $y=0$ gives $f(x) = \phi(x)$ [$\because f(0)=0$]

$$\therefore f(x) + f(y) = f\left(\frac{x+y}{1-xy}\right)$$

Hence, the result.

Section - B

Ques: 5(a) Solve the differential Equation

$$(2y \sin x + 3y^4 \sin x \cos x) dx - (4y^3 \cos^2 x + \cos x) dy = 0$$

Solution:-

Given Equations

$$(2y \sin x + 3y^4 \sin x \cos x) dx - (4y^3 \cos^2 x + \cos x) dy = 0 \quad \text{--- (1)}$$

$$M dx + N dy = 0$$

$$\frac{\partial M}{\partial y} = 2 \sin x + 12y^3 \sin x \cos x$$

$$\frac{\partial N}{\partial x} = 8y^3 \cos x \sin x + \sin x$$

$$\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x} \text{ hence it is not exact.}$$

To make it exact, we need to find I.F.

$$\frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) = \frac{1(\sin x (1 + 4y^3 \cos x))}{-\cos x (4y^3 \cos x + 1)}$$

$$= -\tan x$$

$$\text{I.F} = e^{\int -\tan x dx} = e^{\log \cos x} = \cos x$$

Then; Multiply I.F to equation (1), we get.

$$\Rightarrow (2y \sin x \cos x + 3y^4 \sin x \cos^2 x) dx - (4y^3 \cos^3 x + \cos^2 x) dy = 0$$

$$\Rightarrow (y \sin 2x + \frac{3}{2} y^4 \sin 2x \cos x) dx - (4y^3 \cos x + 1) \frac{(1 + \cos 2x)}{2} dy = 0$$

$$\Rightarrow (y \sin 2x + \frac{3}{2} y^4 \sin 2x \cos x) dx - \left[2y^3 \cos x + 2y^3 \cos x \cos 2x + \frac{1}{2} + \frac{\cos 2x}{2} \right] dy = 0$$

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

$$\Rightarrow \text{Complete solution} = \int_{y=\text{constant}} M dx + \int N dy$$

x-terms not included.

$$\Rightarrow \int_{y=\text{constant}} \left(y \sin 2x + \frac{3}{2} y^4 \sin 2x \cos x \right) dx + \int \frac{1}{2} dy = C$$

$$\Rightarrow -\frac{y \cos 2x}{2} + \frac{3}{4} y^4 \int (\sin 3x + \sin x) dx + \frac{y}{2} = C$$

$$\Rightarrow -\frac{y \cos 2x}{2} + \frac{3}{4} y^4 \left(-\frac{\cos 3x}{3} - \cos x \right) + \frac{y}{2} = C$$

$$\Rightarrow -\left[\frac{y}{2} \cos 2x + \frac{3}{4} y^4 \cos x + \frac{1}{4} y^4 \cos 3x \right] + \frac{y}{2} = C$$

$\Rightarrow$ Multiply whole equation by -4 .

$$\Rightarrow 2y \cos 2x + 3y^4 \cos x + y^4 \cos 3x + (-2y) = C'$$

$$\Rightarrow \boxed{2y \cos 2x + y^4 (\cos 3x + 3 \cos x) - 2y = C'}$$

required solution.

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Ques: 5(b) Determine the complete solution of the differential equation

$$\frac{d^2y}{dx^2} - 4\frac{dy}{dx} + 4y = 3x^2e^{2x}\sin 2x.$$

Solution :- Given;

$$\frac{d^2y}{dx^2} - 4\frac{dy}{dx} + 4y = 3x^2e^{2x}\sin 2x.$$

for the equation, can be written as.

$$y'' - 4y' + 4y = 3x^2e^{2x}\sin 2x. \quad \text{--- (1)}$$

Auxillary equation of $(y'' - 4y' + 4)y = 0$

$$m^2 - 4m + 4 = 0$$

$$(m-2)^2 = 0; \quad m = 2, 2$$

$$\therefore y_c = (C_1 + xC_2)e^{2x} \quad \text{--- (2)}$$

$$\text{for, } y_p = \frac{1}{D^2 - 4D + 4} (3x^2e^{2x}\sin 2x)$$

$$= \frac{1}{(D-2)^2} (3x^2e^{2x}\sin 2x)$$

$$= 3e^{2x} \frac{1}{(D+2-2)^2} x^2 \sin 2x$$

$$= 3e^{2x} \frac{1}{D^2} x^2 \sin 2x. \quad \text{--- (2)}$$

$$\text{let } T = \frac{1}{D^2} x^2 \sin 2x = \frac{1}{D^2} x^2 (\text{I.P. } e^{2xi})$$

$$= \text{I.P. of } \frac{1}{D^2} x^2 \cdot e^{2xi}$$

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

$$= \text{I.P. of } e^{2xi} \frac{1}{(D+2i)^2} x^2$$

$$= \text{I.P. of } e^{2xi} \frac{1}{-4(1+\frac{D}{2i})^2} x^2$$

$$= \text{I.P. of } \frac{e^{2xi}}{-4} \left(1+\frac{D}{2i}\right)^{-2} x^2$$

$$= \text{I.P. of } -\frac{1}{4}(e^{2xi}) \left(1-\frac{D}{i}+\frac{3D^2}{-4}+\dots\right) x^2$$

$$= \text{I.P. of } -\frac{1}{4}(\cos 2x + i \sin 2x) \left[x^2 - \frac{2x}{i} - \frac{3}{4}x \cdot 2\right]$$

$$= \text{I.P. of } -\frac{1}{4}(\cos 2x + i \sin 2x) \left(x^2 + 2xi - \frac{3}{2}\right)$$

$$= \text{I.P. of } -\frac{x}{2} \cos 2x - \left(x^2 - \frac{3}{2}\right) \frac{1}{4} \sin 2x.$$

$$I = -\frac{x}{2} \cos 2x - \frac{x^2}{4} \sin 2x + \frac{3}{8} \sin 2x.$$

$$\therefore y_p = 3e^{2x} \left(-\frac{x}{2} \cos 2x - \frac{x^2}{4} \sin 2x + \frac{3}{8} \sin 2x\right)$$

$$y_p = -3e^{2x} \left[\frac{x}{2} \cos 2x + \frac{x^2}{4} \sin 2x - \frac{3}{8} \sin 2x\right] \quad \text{--- (3)}$$

$$z = y_c + y_p$$

$$z = (C_1 + xC_2)e^{2x} + (-3)e^{2x} \left[\frac{x}{2} \cos 2x + \frac{x^2}{4} \sin 2x - \frac{3}{8} \sin 2x\right]$$

$$z = (C_1 + xC_2)e^{2x} - 3e^{2x} \left[\frac{x}{2} \cos 2x + \frac{x^2}{4} \sin 2x - \frac{3}{8} \sin 2x\right]$$

Required solution.

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Ques: 5(e) Find the directional derivative of the function $xy^2 + yz^2 + zx^2$ along the tangent to the curve $x=t$, $y=t^2$ and $z=t^3$ at the point $(1,1,1)$.

Solution:- let $\phi(x, y, z) = xy^2 + yz^2 + zx^2$

$$\text{Then } \text{grad } \phi = \left(\frac{\partial \phi}{\partial x}\right)\hat{i} + \left(\frac{\partial \phi}{\partial y}\right)\hat{j} + \left(\frac{\partial \phi}{\partial z}\right)\hat{k}$$

$$\text{grad } \phi = (y^2 + 2zx)\hat{i} + (z^2 + 2xy)\hat{j} + (x^2 + 2yz)\hat{k}$$

$$\text{grad } \phi \Big|_{\text{at } (1,1,1)} = 3\hat{i} + 3\hat{j} + 3\hat{k}$$

Also for the curve $x=t$, $y=t^2$ and $z=t^3$ we have, $\frac{dx}{dt} = 1$, $\frac{dy}{dt} = 2t$, $\frac{dz}{dt} = 3t^2$

At point $(1,1,1)$ on the curve $x=t$, $y=t^2$, $z=t^3$ we have $t=1$.

Now vector along the tangent to the above curve at the point (x, y, z)

$$= \left(\frac{dx}{dt}\right)\hat{i} + \left(\frac{dy}{dt}\right)\hat{j} + \left(\frac{dz}{dt}\right)\hat{k}$$

$$= \hat{i} + 2t\hat{j} + 3t^2\hat{k}$$

Putting $t=1$, a vector along the tangent to the curve at the point $(1,1,1)$, we have.

$$= \hat{i} + 2\hat{j} + 3\hat{k}$$

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

If $\hat{a}$ be the unit vector in the direction of this tangent, then

$$\hat{a} = \frac{\hat{i} + 2\hat{j} + 3\hat{k}}{|\hat{i} + 2\hat{j} + 3\hat{k}|} = \frac{\hat{i} + 2\hat{j} + 3\hat{k}}{\sqrt{1+4+9}}$$

$$\hat{a} = \frac{\hat{i} + 2\hat{j} + 3\hat{k}}{\sqrt{14}}$$

$\therefore$ The required directional derivative

$$= \hat{a} \cdot \text{grad } \phi \text{ at } (1, 1, 1)$$

$$= \frac{\hat{i} + 2\hat{j} + 3\hat{k}}{\sqrt{14}} \cdot (3\hat{i} + 3\hat{j} + 3\hat{k})$$


$$= \frac{3+6+9}{\sqrt{14}} = \frac{18}{\sqrt{14}}$$

$$\therefore \boxed{\hat{a} \cdot \text{grad } \phi = \frac{18}{\sqrt{14}}} \text{ required solution}$$

Ques: 6(a) A body consists of a cone and underlying hemisphere. The base of the cone and the top of the hemisphere have same radius a . The whole body rests on a rough horizontal table with hemisphere in contact with the table. Show that the greatest height of the cone, so that the equilibrium may be stable, is $\sqrt{3}a$?

Solution:-

AB is the common base of the hemisphere and the cone and COD is their common axis which must be vertical for equilibrium. The hemisphere touches the table at C. Let H be the height OD of the cone and a be the radius OA or OC or OB of the hemisphere and the cone respectively.



Let G_1 and G_2 be the centres of gravity of the hemisphere and the cone respectively. Then.

$$OG_1 = 3a/8 \text{ and } OG_2 = H/4$$

If h be the height of the centre of gravity of the combined body composed of the hemisphere and the cone above the point of contact 'C', then using the formula

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

$$x = \frac{\omega_1 x_1 + \omega_2 x_2}{\omega_1 + \omega_2}, \text{ we have}$$

$$h = \frac{\frac{1}{3} \pi x^2 H \cdot CG_2 + \frac{2}{3} \pi x^3 CG_1}{\frac{1}{3} \pi x^2 H + \frac{2}{3} \pi x^3} = \frac{\frac{1}{3} \pi x^2 H (x + \frac{1}{4} H) + \frac{2}{3} \pi x^3 \cdot \frac{3}{8} x}{\frac{1}{3} \pi x^2 H + \frac{2}{3} \pi x^3}$$

$$h = \frac{H(x + \frac{1}{4} H) + \frac{5}{4} x^2}{H + 2x}$$

Here; P_1 = radius of curvature at the point of contact C of the upper body, which is spherical = x

P_2 = the radius of curvature of the lower body at the point of contact = ∞

$\therefore$ the equilibrium will be stable if

$$\frac{1}{h} > \frac{1}{P_1} + \frac{1}{P_2} \quad \text{i.e.} \quad \frac{1}{h} > \frac{1}{x} + \frac{1}{\infty} \quad \text{i.e.} \quad \frac{1}{h} > \frac{1}{x}$$

$$\text{i.e.} \quad h < x$$

$$\text{i.e.} \quad \frac{H(x + \frac{1}{4} H) + \frac{5}{4} x^2}{H + 2x} < x$$

$$\text{i.e.} \quad Hx + \frac{1}{4} H^2 + \frac{5}{4} x^2 < Hx + 2x^2$$

$$\text{i.e.} \quad \frac{1}{4} H^2 < \frac{3}{4} x^2 \quad \text{i.e.} \quad H^2 < 3x^2$$

$$\text{i.e.} \quad \boxed{H < x\sqrt{3}} \quad x=a \quad \text{i.e.} \quad \boxed{H < a\sqrt{3}}$$

Hence, the greatest height of the cone consistent with the stable equilibrium of the body is $\sqrt{3}a$ ($\sqrt{3}$ times of the radius) of the hemisphere.

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Ques: 6 (b) Find the circulation of $\vec{F}$ round the curve, where $\vec{F} = (2x+y^2)\hat{i} + (3y-4x)\hat{j}$ and C is the curve $y=x^2$ from $(0,0)$ to $(1,1)$ and the curve $y^2=x$ from $(1,1)$ to $(0,0)$.

Solution:- Here the closed curve 'C' consists of arcs OAP and PBO

Let C_1 denote the arc OAP and C_2 denote arc PBO

Along C_1 , we have $y=x^2$ so that $dy=2x dx$ and x varies from $0 \rightarrow 1$

Along C_2 , we have $x=y^2$

so that $dx=2y dy$ and y varies from 1 to 0 .

Also.


$$F \cdot d\vec{r} = [(2x+y^2)\hat{i} + (3y-4x)\hat{j}] [dx\hat{i} + dy\hat{j}]$$

$$F \cdot d\vec{r} = (2x+y^2)dx + (3y-4x)dy.$$

Now circulation of F round C .

$$\therefore \oint_C F \cdot d\vec{r} = \int_{C_1} \vec{F} \cdot d\vec{r} + \int_{C_2} \vec{F} \cdot d\vec{r}.$$

$$= \int_{C_1} (2x+y^2)dx + (3y-4x)dy + \int_{C_2} (2x+y^2)dx + (3y-4x)dy$$



INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

$$\begin{aligned}
 &= \int_{x=0}^1 (2x+x^4)dx + (3x^2-4x)2xdx \\
 &\quad + \int_{y=1}^0 (2y^2+y^2)2ydy + (3y-4y^2)dy \\
 &= \int_0^1 (2x-8x^2+6x^3+x^4)dx + \int_1^0 (3y-4y^2+6y^3)dy \\
 &= \left[x^2 - \frac{8}{3}x^3 + \frac{3}{2}x^4 + \frac{x^5}{5} \right]_0^1 + \left[\frac{3}{2}y^2 - \frac{4}{3}y^3 + \frac{3}{2}y^4 \right]_1^0 \\
 &= \left[1 - \frac{8}{3} + \frac{3}{2} + \frac{1}{5} \right] - \left[\frac{3}{2} - \frac{4}{3} + \frac{3}{2} \right] \\
 &= \left[\frac{30-80+45+6}{30} \right] - \left[\frac{9-8+9}{6} \right] \\
 &= \frac{1}{30} - \frac{5}{3} = \frac{1-50}{30} = -\frac{49}{30}
 \end{aligned}$$

$$\therefore \oint_C \vec{F} \cdot d\vec{x} = -\frac{49}{30} \quad \text{required solution}$$

-ve sign gives the direction.

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Ques: 6(c)(i) Solve the differential equation

$$\frac{d^2y}{dx^2} + (3\sin x - \cot x) \frac{dy}{dx} + 2y \sin^2 x = e^{-\cos x} \cdot \sin^2 x$$

Solution:- Given that

$$\frac{d^2y}{dx^2} + (3\sin x - \cot x) \frac{dy}{dx} + 2y \sin^2 x = e^{-\cos x} \cdot \sin^2 x \quad \text{--- ①}$$

Clearly it is in the form of

$$\frac{d^2y}{dx^2} + P(x) \frac{dy}{dx} + Q(x)y = R(x),$$

where $P(x) = 3\sin x - \cot x$

$$Q(x) = 2\sin^2 x$$

$$R(x) = e^{-\cos x} \cdot \sin^2 x.$$

let us solve ① by changing the independent variable 'x' to the new independent variable z, where z is a function of 'x'.

Then the transformed equations is -

$$\frac{d^2y}{dz^2} + \frac{\frac{d^2z}{dx^2} + P \frac{dz}{dx}}{\left(\frac{dz}{dx}\right)^2} \cdot \frac{dy}{dx} + \frac{Q(x)}{\left(\frac{dz}{dx}\right)^2} = \frac{R(x)}{\left(\frac{dz}{dx}\right)^2}.$$

let $\frac{Q(x)}{\left(\frac{dz}{dx}\right)^2} = \text{constant}$

$$\Rightarrow \frac{2 \sin^2 x}{\left(\frac{dz}{dx}\right)^2} = 2(\text{say}) \Rightarrow \frac{dz}{dx} = \sin x$$

$$\Rightarrow \boxed{z = \cos x}$$

$\therefore$ from ②

$$\frac{d^2y}{dz^2} + \frac{-\cos x + (3\sin x - \cot x)(-\sin x)}{\sin^2 x} \left(\frac{dy}{dx}\right) + 2y = \frac{e^{-\cos x} \cdot \sin^2 x}{\sin^2 x}$$

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

$$\Rightarrow \frac{d^2y}{dz^2} + \frac{-\cos x - 3 \sin^2 x + \cos x}{\sin^2 x} \left(\frac{dy}{dx} \right) + 2y = e^{-\cos x}$$

$$\Rightarrow \frac{d^2y}{dz^2} - 3 \frac{dy}{dz} + 2y = e^{-z}$$

$$(D^2 - 3D + 2)y = e^{-z} \quad ; \quad D = \frac{d}{dz} \quad \text{--- (3)}$$

Its A.E. is $m^2 - 3m + 2 = 0$
 $(m-2)(m-1) = 0$
 $m = 1, 2.$

$$\therefore y_c(z) = ae^z + be^{2z} \quad \text{--- (4)}$$

and we have.

$$y_p(z) = \frac{1}{D^2 - 3D + 2} e^{-z} = \frac{1}{1+3+2} e^{-z}$$

$$y_p(z) = \frac{1}{6} e^{-z}.$$

$\therefore$ The general solution of (1) is
 given by $y(z) = y_c(z) + y_p(z)$

$$y(z) = ae^z + be^{2z} + \frac{1}{6} e^{-z}$$

$$\Rightarrow y(x) = ae^{\cos x} + be^{2\cos x} + \frac{1}{6} e^{-\cos x}.$$

which is the required general solution
 of the given equation.

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Ques:- 6(c) ii) Find the Laplace transforms of $t^{-1/2}$ and $t^{1/2}$. Prove that the Laplace transform of $t^{n+1/2}$, where $n \in \mathbb{N}$, is

$$\frac{\Gamma\left(n+1+\frac{1}{2}\right)}{s^{n+1+\frac{1}{2}}}.$$

Solution :-

(a) $f(t) = t^{-1/2}$

$$L f(t) = \int_0^{\infty} e^{-st} \frac{1}{\sqrt{t}} dt$$

Now put $st = p \Rightarrow s dt = dp$

$$\therefore dt = \frac{dp}{s}$$

$$\Rightarrow \int_0^{\infty} e^{-p} \frac{\sqrt{s}}{\sqrt{p}} \cdot \frac{1}{s} dp = \frac{1}{\sqrt{s}} \int_0^{\infty} e^{-p} \cdot p^{-1/2} dp.$$

$$\Rightarrow \frac{1}{\sqrt{s}} \cdot \sqrt{\frac{\pi}{2}} = \frac{\sqrt{\pi}}{\sqrt{s}} = \underline{\underline{\frac{\sqrt{\pi}}{\sqrt{s}}}} \quad \left\{ \begin{array}{l} \text{By gamma} \\ \text{function} \end{array} \right\}$$

(b) $f(t) = t^{1/2}$

$$L f(t) = \int_0^{\infty} e^{-st} \sqrt{t} dt$$

put $st = p \Rightarrow s dt = dp \Rightarrow dt = \frac{dp}{s}$

$$\Rightarrow \int_0^{\infty} e^{-p} \frac{\sqrt{p}}{\sqrt{s}} \cdot \frac{dp}{s} = s^{-3/2} \int_0^{\infty} e^{-p} p^{1/2} dp.$$

$$= s^{-3/2} \int_0^{\infty} e^{-p} \cdot p^{3/2-1} dp.$$

(By gamma function)

$$L(f(t)) = \frac{1}{2s} \sqrt{\frac{\pi}{s}}$$

Required Solution.

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

© $f(t) = t^n$

$$\therefore f(s) = L(f(t)) = \lim_{A \rightarrow \infty} \int_0^A e^{-st} \cdot t^n dt$$

$$= \lim_{A \rightarrow \infty} \left\{ t^n \cdot \frac{e^{-st}}{-s} \Big|_0^A - \int_0^A \frac{xt^{n-1} \cdot e^{-st}}{-s} dt \right\}$$

$$= 0 + \frac{n}{s} \lim_{A \rightarrow \infty} \int_0^A e^{-st} t^{n-1} dt$$

$$L\{t^n\} = \frac{n}{s} L\{t^{n-1}\}$$

So recursive solution

$$L\{t^n\} = \frac{n}{s} \{L\{t^{n-1}\}\}, \forall n$$

So, $L\{t^{n-1}\} = \frac{n-1}{s} L\{t^{n-2}\}$

$$L\{t^{n-2}\} = \frac{n-2}{s} L\{t^{n-3}\}$$

by Mathematical Induction, we get

$$L\{t^n\} = \frac{n}{s} \cdot \frac{(n-1)}{s} \cdot \frac{(n-2)}{s} \cdots \frac{1}{s} L\{1\}$$

$$= \frac{n!}{s^n} \cdot \frac{1}{s} = \frac{n!}{s^{n+1}}$$

$\therefore$ from above formula

$$L\{t^{n+1/2}\} = \frac{\Gamma(n+1+1/2)}{s^{n+1+1/2}}$$

Hence the result.

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Ques: 7(a) Find the linearly independent solutions of the corresponding homogeneous differential equation of the equation $x^2 y'' - 2xy' + 2y = x^3 \sin x$ and then find the general solution of the given equation by the method of variation of parameters.

Solution: Given

$$x^2 y'' - 2xy' + 2y = x^3 \sin x \quad \text{--- (1)}$$

Divide equation (1) by x^2 .

$$y'' - \frac{2}{x} y' + \frac{2}{x^2} y = x \sin x.$$

[$\because R = x \sin x$]

$$y'' - \frac{2}{x} y' + \frac{2}{x^2} y = 0 \quad \text{homogeneous equation}$$

$$x^2 y'' - 2xy' + 2y = 0$$

$$(x^2 y'' - 2xy' + 2)y = 0$$

$$(x^2 D^2 - 2xD + 2)y = 0 \quad [\because D \equiv \frac{d}{dx}]$$

put $x = e^z$ and $D_1 \equiv \frac{d}{dz}$

Then, we get

$$(D_1(D_1 - 1) - 2(D_1) + 2)y = 0$$

$$\Rightarrow (D_1^2 - 3D_1 + 2)y = 0$$

$$(D_1 - 2)(D_1 - 1) = 0$$

$$D = 2, 1.$$

$$y_c = C_1 e + C_2 e^2$$

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

In terms of x .

$$y_c = C_1 x + C_2 x^2 \quad [\because x = e^z]$$

where $u = x \quad v = x^2$
 $u' = 1 \quad v' = 2x$.

$$W = \begin{vmatrix} u & v \\ u' & v' \end{vmatrix} = \begin{vmatrix} x & x^2 \\ 1 & 2x \end{vmatrix} = 2x^2 - x^2 = x^2 \neq 0$$

for P.I. = $A \cdot u + B \cdot v$.

where $A = -\int \frac{vR}{W} dx$ and $B = \int \frac{uR}{W} dx$.

$$A = -\int \frac{x^2 \cdot x \sin x}{x^2} dx$$

$$A = -(x \cos x + \sin x) = x \cos x - \sin x + C_1$$

$$B = \int \frac{x \cdot x \sin x}{x^2} dx = \int \sin x dx$$

$$B = -\cos x + C_2$$

$$P.I. = x \cdot [x \cos x - \sin x] + x^2 [-\cos x]$$

$$P.I. = x^2 \cos x - x \sin x + (-x^2 \cos x)$$

$$\boxed{P.I. = -x \sin x}$$

$$z = y_c + y_p$$

$$\boxed{z = C_1 x + C_2 x^2 - x \sin x}$$

Required general solution.

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Ques: 7(b) Find the radius of curvature and radius of torsion of the helix $x = a \cos u$, $y = a \sin u$, $z = a \tan \alpha$.

Solution : Given; $x = a \cos u$ $y = a \sin u$
 $z = a \tan \alpha$

$$\mathbf{r} = x\hat{i} + y\hat{j} + z\hat{k}$$

$$\mathbf{r} = a \cos u \hat{i} + a \sin u \hat{j} + a \tan \alpha \hat{k} \quad \text{--- (1)}$$

$$\kappa = \frac{\left| \frac{d\mathbf{r}}{du} \times \frac{d^2\mathbf{r}}{du^2} \right|}{\left| \frac{d\mathbf{r}}{du} \right|^3} \quad \text{--- (2)}$$

$$\tau = \frac{\left| \frac{d\mathbf{r}}{du} \cdot \frac{d^2\mathbf{r}}{du^2} \cdot \frac{d^3\mathbf{r}}{du^3} \right|}{\left| \frac{d\mathbf{r}}{du} \times \frac{d^2\mathbf{r}}{du^2} \right|^2} \quad \text{--- (3)}$$

from (1); differentiate w.r.t 'u' -

$$\frac{d\mathbf{r}}{du} = -a \sin u \hat{i} + a \cos u \hat{j} + a \tan \alpha \hat{k}$$

$$\frac{d^2\mathbf{r}}{du^2} = -a \cos u \hat{i} - a \sin u \hat{j} + 0 \hat{k}$$

$$\frac{d^3\mathbf{r}}{du^3} = +a \sin u \hat{i} - a \cos u \hat{j}$$

$$\begin{aligned} \left| \frac{d\mathbf{r}}{du} \right| &= \sqrt{(-a \sin u)^2 + (a \cos u)^2 + (a \tan \alpha)^2} \\ &= \sqrt{a^2(\sin^2 u + \cos^2 u + \tan^2 \alpha)} \\ &= a \sqrt{1 + \tan^2 \alpha} = a \sqrt{\sec^2 \alpha} \end{aligned}$$

$$\boxed{\left| \frac{d\mathbf{r}}{du} \right| = a \sec \alpha}$$

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

$$\begin{aligned} \frac{dx}{du} \times \frac{d^2x}{du^2} &= \begin{vmatrix} i & j & k \\ -a \sin u & a \cos u & a \tan \alpha \\ -a \cos u & -a \sin u & 0 \end{vmatrix} \\ &= a^2 \begin{vmatrix} i & j & k \\ -\sin u & \cos u & \tan \alpha \\ -\cos u & -\sin u & 0 \end{vmatrix} \\ &= a^2 [i(-\sin u \cdot \tan \alpha) - \cos u \tan \alpha \hat{j} + \hat{k}] \end{aligned}$$

$$\begin{aligned} \left| \frac{dx}{du} \times \frac{d^2x}{du^2} \right| &= \sqrt{a^4 (\sin^2 u \tan^2 \alpha + \cos^2 u \tan^2 \alpha) + a^4} \\ &= a^2 \sqrt{\tan^2 \alpha (\sin^2 u + \cos^2 u) + 1} \\ &= a^2 \sqrt{\tan^2 \alpha + 1} = a^2 \sec \alpha. \end{aligned}$$

$$\begin{aligned} \text{Curvature} = K &= \frac{a^2 \sec \alpha}{(a \sec \alpha)^3} = \frac{a^2 \sec \alpha}{a^3 \cdot \sec^3 \alpha} \\ K &= \frac{1}{a} \cdot \frac{1}{\sec^2 \alpha} = \frac{1}{a} \cos^2 \alpha. \end{aligned}$$

$$\boxed{\text{Radius of curvature} = \frac{1}{K} = a \sec^2 \alpha.}$$

$$\begin{aligned} \left[\frac{dx}{du} \frac{d^2x}{du^2} \frac{d^3x}{du^3} \right] &= \begin{vmatrix} a \sin u & -a \cos u & 0 \\ -a \sin u & a \cos u & a \tan \alpha \\ a \cos u & a \sin u & 0 \end{vmatrix} \\ &= a^3 \begin{vmatrix} \sin u & -\cos u & 0 \\ -\sin u & \cos u & \tan \alpha \\ \cos u & \sin u & 0 \end{vmatrix} \end{aligned}$$

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

$$R_2 \rightarrow R_2 + R_1$$

$$= a^3 \begin{vmatrix} \sin u & -\cos u & 0 \\ 0 & 0 & \tan \alpha \\ \cos u & \sin u & 0 \end{vmatrix}$$

$$= a^3 [\sin u [-\sin u \tan \alpha] + \cos u [-\cos u \tan \alpha] + 0]$$

$$= a^3 [-[\sin^2 u \tan \alpha + \cos^2 u \tan \alpha]]$$

$$= -a^3 [\tan \alpha [\sin^2 u + \cos^2 u]]$$

$$= -a^3 \tan \alpha \quad [\because \sin^2 u + \cos^2 u = 1]$$

$$\tau = \frac{|-a^3 \tan \alpha|}{(a^2 \cdot \sec \alpha)^2} = \frac{a^3 \tan \alpha}{a^4 \cdot \sec^2 \alpha}$$

$$\tau = \frac{1}{a} \cdot \frac{1}{2} \cdot \sin 2\alpha = \frac{1}{2a} \cdot \sin 2\alpha$$

$$\therefore \text{radius of Torsion} = \frac{1}{\tau} = \frac{a \sec^2 \alpha}{\tan \alpha}$$

$$\text{radius of Torsion} = \frac{2a}{\sin 2\alpha}$$

required solution.

Ques: 8(a) Obtain the singular solution of the differential equation

$$\left(\frac{dy}{dx}\right)^2 \left(\frac{y}{x}\right)^2 \cot^2 \alpha - 2 \left(\frac{dy}{dx}\right) \left(\frac{y}{x}\right) + \left(\frac{y}{x}\right)^2 \operatorname{cosec}^2 \alpha = 1$$

Also find the complete primitive of the given differential equation. Give the geometrical interpretations of the complete primitive and singular solution.

Solution :-

Given ; $\left(\frac{dy}{dx}\right)^2 \left(\frac{y}{x}\right)^2 \cot^2 \alpha - 2 \left(\frac{dy}{dx}\right) \left(\frac{y}{x}\right) + \left(\frac{y}{x}\right)^2 \operatorname{cosec}^2 \alpha = 1$

which can be rewritten as.

$$p^2 y^2 \cot^2 \alpha - 2pxy + y^2 \operatorname{cosec}^2 \alpha = x^2.$$

$$p^2 y^2 \frac{\cos^2 \alpha}{\sin^2 \alpha} - 2pxy + y^2 \frac{1}{\sin^2 \alpha} = x^2$$

$$p^2 y^2 \cos^2 \alpha - 2pxy \sin^2 \alpha + y^2 = x^2 \sin^2 \alpha.$$

$$\Rightarrow p^2 y^2 \cos^2 \alpha - 2pxy \sin^2 \alpha + y^2 - x^2 \sin^2 \alpha = 0 \quad \text{--- (A)}$$

$$\Rightarrow (py)^2 - (2py)x \tan^2 \alpha + (y^2 \sec^2 \alpha - x^2 \tan^2 \alpha) = 0$$

$$\therefore py = \frac{2x \tan^2 \alpha \pm \sqrt{4x^2 \tan^4 \alpha - 4(y^2 \sec^2 \alpha - x^2 \tan^2 \alpha)}}{2}.$$

$$\text{or } py = x \tan^2 \alpha \pm \sqrt{x^2 \tan^2 \alpha (\tan^2 \alpha + 1) - y^2 \sec^2 \alpha}$$

$$py = x \tan^2 \alpha \pm \sec \alpha \sqrt{x^2 \tan^2 \alpha - y^2}$$

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

$$y dy - x \tan^2 \alpha dx = \pm \sec \alpha \sqrt{(x^2 \tan^2 \alpha - y^2)} dx.$$

$$\Rightarrow \pm \frac{x \tan^2 \alpha dx - y dy}{\sqrt{(x^2 \tan^2 \alpha - y^2)}} = -\sec \alpha dx$$

Integrating, $\pm \sqrt{x^2 \tan^2 \alpha - y^2} = C - x \sec \alpha$

Squaring; $x^2 \tan^2 \alpha - y^2 = C^2 - 2Cx \sec \alpha + x^2 \sec^2 \alpha$

$$\Rightarrow x^2(\tan^2 \alpha - \sec^2 \alpha) - y^2 = C^2 - 2Cx \sec \alpha$$

$$\Rightarrow -x^2 - y^2 = C^2 - 2Cx \sec \alpha \quad [\because 1 = \tan^2 \alpha - \sec^2 \alpha]$$

$$\Rightarrow x^2 + y^2 - 2Cx \sec \alpha + C^2 = 0 \quad \text{--- (B)}$$

From (A)

$$p^2 y^2 \cos^2 \alpha - 2pxy \sin^2 \alpha + y^2 - x^2 \sin^2 \alpha = 0$$

Since the given equation is quadratic in p, the p-discriminant relation is -

$$4x^2 y^2 \sin^4 \alpha - 4y^2 \cos^2 \alpha (y^2 - x^2 \sin^2 \alpha) = 0$$

$$4y^2 [x^2 \sin^2 \alpha (\sin^2 \alpha + \cos^2 \alpha) - y^2 \cos^2 \alpha] = 0$$

$$y^2 [x^2 \sin^2 \alpha - y^2 \cos^2 \alpha] = 0$$

$$y^2 \cos^2 \alpha (x^2 \tan^2 \alpha - y^2) = 0 \quad \text{--- (C)}$$

Now, the general solution of (A) is

$$C^2 - 2Cx \sec \alpha + x^2 + y^2 = 0 \quad \text{--- (D)}$$

$\therefore$ The c-discriminant relation is

$$4x^2 \sec^2 \alpha - 4(x^2 + y^2) = 0$$

$$x^2(\sec^2\alpha - 1) - y^2 = 0$$

$$x^2 \tan^2\alpha - y^2 = 0$$

$$\therefore (x \tan\alpha - y)(x \tan\alpha + y) = 0 \quad \text{--- (D)}$$

$$\text{So; } x \tan\alpha - y = 0 \quad x \tan\alpha + y = 0$$

i.e. the lines $y = \pm x \tan\alpha$ are singular solution (envelopes) and $y=0$ is a tac-locus.

Now the general solution (C) represents a family of circles all having their centres on x-axis. The circles of the system touch one another on x-axis and $y=0$ i.e. x-axis passes through the points of contact of non-consecutive circles which touch on x-axis. The family of circles is being touched by $y = \pm x \tan\alpha$, which are equally inclined to the line of centres of the circles i.e. x-axis and pass through the origin.

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Ques: 8(b) Prove that the path of a planet, which is moving so that its acceleration is always directed to a fixed point (star) and is equal to $\frac{\mu}{(\text{distance})^2}$, is a conic section. Find the conditions under which the path becomes (i) ellipse (ii) parabola and (iii) hyperbola.

Solution:-

Here the force is always directed to a fixed point (star), so it is a case of central orbit.

Also given that the central acceleration $p = \frac{\mu}{r^2}$

$r = \text{distance}$

The differential equation of the path (in pedal form) is

$$\frac{h^2}{p^3} \frac{dp}{dr} = p = \frac{\mu}{r^2}$$

Multiplying both sides by -2 , we have

$$-\frac{2h^2}{p^3} dp = -\frac{2\mu}{r^2} dr$$

Integrating, we have

$$v^2 = \frac{h^2}{p^2} = \frac{2\mu}{r} + B \quad \text{--- (1)}$$

where 'B' is a constant

[Note that in a central orbit, $v = h/p$]

We know that referred to the focus as pole the pedal equations of ellipse, parabola and hyperbola (that branch which is nearer to focus taken as pole) are

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

$$\frac{b^2}{p^2} = \frac{2a}{r} - 1, \quad p^2 = ar \quad \text{and} \quad \frac{b^2}{p^2} = \frac{2a}{r} + 1$$

respectively, where
 in the case of ellipse $2a$ and $2b$ are the lengths of major and minor axes,
 in the case of parabola $4a$ is the length of latus rectum, and in the case of hyperbola $2a$ and $2b$ are the lengths of transverse and conjugate axes.

Now, since the equation (1) can be any of the above three forms, three cases arise here.

Case I: Elliptic Path

Comparing (1) with $\frac{b^2}{p^2} = \frac{2a}{r} - 1$, the pedal equation of the ellipse, we have.

$$\frac{h^2}{b^2} = \frac{\mu}{a} = \frac{B}{-1}.$$

$$\therefore h^2 = \frac{\mu b^2}{a} \quad \text{and} \quad B = -\frac{\mu}{a}.$$

Substituting in (1), for elliptical path, we have.

$$v^2 = \frac{2\mu}{r} - \frac{\mu}{a} = \mu \left(\frac{2}{r} - \frac{1}{a} \right)$$

obviously here $\boxed{v^2 < \frac{2\mu}{r}}$

Case II: Parabolic Path:- Comparing (1) with $p^2 = ar$, the pedal equation of a parabola, we have

$$\frac{h^2}{1} = \frac{2\mu}{4/a} = \frac{B}{0}$$

$$\therefore h^2 = 2\mu a \quad \text{and} \quad B = 0.$$

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Substituting in (1), for parabolic path we have

$$\boxed{v^2 = \frac{2\mu}{r}}$$

Case II: Hyperbolic path: Comparing (1) with

$\frac{b^2}{\rho^2} = \frac{2a}{r} + 1$ the pedal equation of a hyperbola, we have

$$\frac{h^2}{b^2} = \frac{\mu}{a} = \frac{B}{1}$$

$$\therefore h^2 = \frac{\mu b^2}{a} \text{ and } B = \mu/a.$$

Substituting in (1), for hyperbolic path, we have

$$v^2 = \mu \left(\frac{2}{r} + \frac{1}{a} \right)$$

obviously here $\boxed{v^2 > \frac{2\mu}{r}}$

Thus from the above three cases, we conclude that the equation (1) always represents a conic section whose focus is at the centre of force. Further the path of the particle is an ellipse, parabola and hyperbola according as B is -ve, zero and positive. The sign of the value of the constant B depends upon the magnitude of the velocity of the particle at any point. we have found that

if (i) $v^2 = \mu \left(\frac{2}{r} - \frac{1}{a} \right)$ or $v^2 < \frac{2\mu}{r}$

then path is elliptic,

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

if $v^2 = \frac{2\mu}{r}$, then the path is parabolic

if $v^2 > \frac{2\mu}{r}$ or $v^2 = \frac{2\mu}{r} + \frac{\mu}{a}$, then the path is hyperbolic.

It is to be noted that in each of the three cases magnitude of the velocity at any point is independent of the direction of the velocity at that time.

Also, we have found that -

$$h^2 = \mu b^2/a = \mu l \quad \text{in case of elliptic path}$$

$$h^2 = 2\mu a = \mu l \quad \text{in case of parabolic path}$$

$$h^2 = \mu b^2/a = \mu l \quad \text{in case of hyperbolic path.}$$

Thus in all the three cases.

$h = \sqrt{\mu l}$, where 'l' is the length of the semi-latus rectum.

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Ques: 8(C)(i) State Gauss divergence theorem. Verify this theorem for $\vec{F} = 4x\hat{i} - 2y^2\hat{j} + z^2\hat{k}$, taken over the region bounded by $x^2 + y^2 = 4$, $z = 0$ and $z = 3$.

Solution:-

Gauss Divergence theorem states that if V is the volume bounded by a closed surface S and $\vec{A}$ is a vector function of position with continuous derivatives, then

$$\boxed{\iiint_V \nabla \cdot \vec{A} \, dv = \iint_S \vec{A} \cdot \hat{n} \, ds}$$

where $\hat{n}$ is the positive (outward drawn) normal to S .

Now; $\vec{F} = 4x\hat{i} - 2y^2\hat{j} + z^2\hat{k}$

taken over the region bounded by $x^2 + y^2 = 4$, $z = 0$ and $z = 3$.

$\therefore$ By Gauss divergence theorem

$$\iiint_V \nabla \cdot \vec{F} \, dv = \iint_S \vec{F} \cdot \hat{n} \, ds$$

$$\text{volume integral} = \iiint_V \nabla \cdot \vec{F} \, dv$$

$$= \iiint_V \left[\frac{\partial}{\partial x} (4x) + \frac{\partial}{\partial y} (-2y^2) + \frac{\partial}{\partial z} (z^2) \right] dv$$

$$= \iiint_V (4 - 4y + 2z) dv$$

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

$$= \int_{x=-2}^2 \int_{y=-\sqrt{4-x^2}}^{\sqrt{4-x^2}} \int_{z=0}^3 (4-4y+2z) dz dy dx$$

$$= \int_{x=-2}^2 \int_{-\sqrt{4-x^2}}^{\sqrt{4-x^2}} [4z - 4yz + z^2]_0^3 dy dx \quad \left[\begin{array}{l} \because \text{radius of circle} \\ x^2 + y^2 = 4 \quad x=2 \\ y = \pm \sqrt{4-x^2} \end{array} \right]$$

$$= \int_{x=-2}^2 \int_{-\sqrt{4-x^2}}^{\sqrt{4-x^2}} [12 - 12y + 9] dy dx$$

using polar coordinates

$$x = r \cos \theta$$

$$y = r \sin \theta$$

$$dx = r(-\sin \theta) d\theta$$

$$dy = r \cos \theta d\theta$$

$$\Rightarrow dx dy = r d\theta dr$$

$$\int_0^{2\pi} \int_0^2 (21 - 12r \sin \theta) r dr d\theta$$

$$= \int_0^{2\pi} \int_0^2 (21r - r^2 \sin \theta) dr d\theta$$


$$= \int_0^{2\pi} \left[\frac{21r^2}{2} - \frac{r^3}{3} \sin \theta \right]_0^2 d\theta$$

$$= \int_0^{2\pi} \left[42 - \frac{8}{3} \sin \theta \right] d\theta = \left[42\theta + \frac{8}{3} \cos \theta \right]_0^{2\pi}$$

$$= \left[84\pi + \left[\frac{8}{3}(1) - \frac{8}{3}(1) \right] \right] = 84\pi$$

$$\Rightarrow \boxed{\iiint_V \nabla \cdot \vec{F} dv = 84\pi}$$

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna



The surface S of the cylinder consists of a base $S_1 (z=0)$, the top $S_2 (z=3)$ and the convex portion $S_3 (x^2+y^2=4)$. Then.

$$\text{Surface Integral} = \iint_S \vec{F} \cdot \hat{n} ds = \iint_{S_1} \vec{F} \cdot \hat{n} ds_1 + \iint_{S_2} \vec{F} \cdot \hat{n} ds_2 + \iint_{S_3} \vec{F} \cdot \hat{n} ds_3$$

On $S_1 (z=0)$; $\hat{n} = -\hat{k}$ $\vec{F} = 4x\hat{i} - 2y^2\hat{j}$

$$\vec{F} \cdot \hat{n} = 0$$

$$\therefore \iint_{S_1} \vec{F} \cdot \hat{n} ds_1 = 0.$$

On $S_2 (z=3)$; $\hat{n} = \hat{k}$ $\vec{F} = 4x\hat{i} - 2y^2\hat{j} + 9\hat{k}$

$$\vec{F} \cdot \hat{n} = 9, \text{ so that}$$

$$\begin{aligned} \iint_{S_2} \vec{F} \cdot \hat{n} ds_2 &= \iint_{S_2} 9 ds_2 = 9 \cdot \pi r^2 \\ &= 9\pi \cdot (2)^2 \quad [\because r=2] \end{aligned}$$

$$\boxed{\iint_{S_2} \vec{F} \cdot \hat{n} ds_2 = 36\pi}$$

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

On $S_3(x^2+y^2=4)$:

A perpendicular to $x^2+y^2=4$ has the direction

$$\nabla(x^2+y^2) = 2x\hat{i} + 2y\hat{j}$$

Then a unit normal is $\hat{n} = \frac{2x\hat{i} + 2y\hat{j}}{\sqrt{4x^2 + 4y^2}}$

$$\hat{n} = \frac{2(x\hat{i} + y\hat{j})}{2\sqrt{x^2 + y^2}} = \frac{x\hat{i} + y\hat{j}}{2} \quad [\because x^2 + y^2 = 4]$$

$$\vec{F} \cdot \hat{n} = (4x\hat{i} - 2y^2\hat{j} + z^2\hat{k}) \left(\frac{x\hat{i} + y\hat{j}}{2} \right)$$

$$\boxed{\vec{F} \cdot \hat{n} = 2x^2 - y^3}$$

$$\therefore \iint_{S_3} \vec{F} \cdot \hat{n} dS = \int_{\theta=0}^{2\pi} \int_{z=0}^3 [2(2\cos\theta)^2 - (2\sin\theta)^3] 2 dz d\theta$$

$$= \int_{\theta=0}^{2\pi} (48\cos^2\theta - 48\sin^3\theta) d\theta$$

$$= \int_{\theta=0}^{2\pi} 48\cos^2\theta d\theta = 48\pi.$$

$$\therefore \text{Then the surface integral} = 0 + 36\pi + 48\pi = 84\pi.$$

$$\therefore \boxed{\iint_S \vec{F} \cdot \hat{n} dS = \iiint_V \vec{F} dV = 84\pi}$$

Hence verified.

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Ques: 8(c) ii) Evaluate by stoke's theorem $\oint_C e^x dx + 2y dy - dz$,
 where C is the curve $x^2 + y^2 = 4, z = 2$.

Solution:-

$$\vec{F} = e^x dx + 2y dy - dz \quad \& \quad \oint_C \vec{F}$$

C is the curve $x^2 + y^2 = 4; z = 2$

Using stokes theorem

$$\oint_C \vec{F} \cdot d\vec{r} = \iint (\nabla \times \vec{F}) \cdot \hat{n} ds$$

$$\nabla \times \vec{F} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \partial/\partial x & \partial/\partial y & \partial/\partial z \\ e^x & 2y & -1 \end{vmatrix}$$

$$\nabla \times \vec{F} = \hat{i}(0) - \hat{j}(0) + \hat{k}(0)$$

$$\nabla \times \vec{F} = 0$$

$$\therefore \iint (\nabla \times \vec{F}) \cdot \hat{n} ds = 0$$

$$\therefore \boxed{\oint_C e^x dx + 2y dy - dz = \iint (\nabla \times \vec{F}) \cdot \hat{n} ds = 0}$$

which is required result